University of Mumbai No.UG/ of 2007

CIRCULAR:

In partial modification of this office Circular No.UG/476 of 2006 dated 1st December, 2006, the Principals of the affiliated colleges in Science are hereby informed that the Appendix sent alongwith the above mentioned circular is treated as cancelled and that the correct Appendix of the guidelines for practical in the subject of Statistics at the TY.B.Sc. is enclosed herewith for your information.

Mumbai 400 032 19th January, 2007

To,

The Principals of the affiliated colleges in Science.

A.C.4.7/18.08.06

No.UG/ 22-A of 2007

January, 2007

for REGISTR

for REGIS

Copy forwarded with compliments to :-

1) the Dean, Faculty of Science,

2) the Chairman, Board of Studies in Statistics.

Copy to:

The Director, Board of College and University Development, Deputy Registrar (Eligibility and Migration), the Director of students Welfare, the Personal Assistants to the Vice-Chancellor, the Pro-Vice-Chancellor, the Registrar and the Assistant Registrar Administrative sub-centre, Ratnagiri, for information.

The O.S.D. and Controller of Examinations (10 copies), the F. & A.O. (Accounts Section), Fort (2 copies), Record Section (5 copies), Publication Section (5 copies), D.R., Enrolment (3 copies), D.R. (Statistical Unit), D.R., Accounts Section, Vidyanagari, (2 copies), the Director, University Computer Centre (I.D.E. Bldg.), Vidyanagari, (2 copies), D.R. (Aff. Section) (2 copies), A.R., A.A. Unit (2 copies) and A.R. (Executive Authorities Unit) (2 copies), they are requested to treat this as action taken report on the concerned resolution adopted by the Academic Council/Management Council referred to in the above Circular and that no separate ATR. will be sent in this connection. A.R., CONCOL (1 copy), BUCTU (2 copies), Incharge, C.C.F. (1 copy), D.R., I.D.E. (10 copies), D.R., P.R.O., (2 copies), D.R. (Special Cell), Supdt., Thesis Section (2 copies), Supdt., P.G. Section (2 copies), Secretary, MUASA (1 copy), Dy. Acett. (Unit V) (1 copy), Receptionist (1 copy), Telephone Operator (1 copy).

Nsk/Cir.2/170107

UNIVERSITY OF MUMBAI



GUIDELINES FOR PRACTICAL. IN THE SUBJECT OF **STATISTICS** AT THE T.Y.B.SC.

UNIT-VI (Sections I & II) UNIT-VII (Sections I & II) UNIT-VIII (Sections I & II) UNIT-IX (Sections I & II)

With effect from the academic year 2005-2007)

2007-2008

T.Y.B.Sc. STATISTICS PRACTICALS INDEX

		The state of the s
it Sec	Prac No	little
1 .50	0 1 1	Prohability
	6.12	Probability Generating function
	6.1.3	Stochastic Processes
	614	Queueing Theory-I
1 .	615	Queueing Theory-II
1-0	6.2.1	Chebychev's Inequality
11	6.2.2	Order Statistics
	6.2.3	Bivariate Discrete Distributions
	6.2.4	Bivariate Continuous Distributions
1	6.2.5	Significance of Correlation Coefficient
1	6.2.6	Multiple Partial Correlation and D
1	0.2.0	Multiple, Partial Correlation and Regression
1	7.1.1	Method of Maximum Likelihood, Method of Moments and
1	,,,,,,	Method of Minimum Chi-equare
	7.1.2	Method of Maximum Likelihood and Method of Moments
	7.1.3	Bayes' Estimation
	7.1.4	Interval Estimation
	7.1.5	Linear Models
	/.1.5	Lineal Models
П	7.2.1	Testing of Hypothesis-I
	7.2.2	Testing of Hypothesis-II
	7.2.3	Likelihood Ratio Test
	7.2.4	Non-parametric Test-I
	7.2.5	Non-parametric Test-II
	7.2.6	Sequential Probability Ratio Test (S.P.R.T.)
	/12.0	sequential Probability Ratio Test (S.F.R. 1.)
I	8.1.1	Genetics-I
	8.1.2	Genetics-II
	8.1.3	Genetics-III
	8.1.4	Epidemics-I
	8.1.5	Epidemics-II
	8.1.6	
	8.1.7	Bio-assays-I
		Bio-assays-II
	8.1.8	Bio-assays-III
П	8.2.1	Impentory Models I
	8.2.2	Inventory Models-I
	000	Inventory Models-II
	8.2.4	Replacement Theory
		Decision Theory
	8.2.5	Theory of Games
-		
1		

01.1	Vital Statistics
012	Life Tables-I
	Life Tables-II
	Armuitics-I
	Annuitics-II
	Premiums and Assurances
9.1.7	Life Annuities and Temporary Annuities
021	Reliability
-	Time Series
	Simulation
	Mathematical Economics-I
	Mathematical Economics-II
	Input-Output Models
9.2.0	The surface states
	9.1.1 9.1.2 9.1.3 9.1.4 9.1.5 9.1.6 9.1.7 9.2.1 9.2.2 9.2.3 9.2.4 9.2.5 9.2.6

T.Y.B.Sc. Practical No.6.1.1

PROBABILITY

In 'Maharaja' hotel there are 10 rooms, each having capacity of maximum 40 persons. In 'Manaraja Union on the Sunday 40 passengers arrived at the hotel and manager places them at on one in 10 rooms. What is the probability that a) no room is empty?

b) exactly 7 rooms remain empty?

b) exactly (states) b) exactly (states) b) exactly (states) exactly (state a) all passengers leave on same floor?

b) each passenger leaves on different floor?

c) on at least 3 floors, no passenger leaves?

Five different letters are distributed at random in 5 self addressed envelopes. Find the probability that

a) exactly 2 letters

b) at least 3 letters and

c) none

are placed in correct envelope.

- A student is asked to guess an unknown order of 10 distinct cards. What is the probability of exactly 4 correct guesses if he calls each card only once at random? What is the probability that all his guesses an wrong?
- A pack of cards consists of 5 identical series, each containing 10 cards numbered 1 to 10. A random sample of size 15 is drawn from the pack
 - a) without replacement

b) with replacement.

Calculate the probability that each number is represented in the sample.

- Twenty pairs of shoes are in a closet. Six shoes are selected at random. Find the probability that there will be
 - a) at least one pair
 - b) exactly 2 pairs

among the 6 shoes selected.

- 7) In a game of bridge, what is the chance that
 - a) exactly 2 players
 - b) at least 3 players

get complete suit?

1) In a lottery, 2 tickets are drawn at a time out of the total number of 5 nekets numbered I to 5 and returned before the next drawing is made. Find the chance that in 3 drawings, each of the numbers 1,2..... 5 will appear at least once

1) The probability distribution function of a random variable X is
$$p(x) = (1/6) (5/6)^{x-1}$$
; $x = 1, 2, 3, ...$; otherwise

a) probability generating function of X.

b) E(X) and V(X).

The probability of error in the transmission of a bit over a communication channel is $p = 10^{-4}$ Let X denotes the number of errors in transmitting a block of 1000 bits. Find a) probability generating function of X.

b) variance of the distribution of X.

A random variable Y is defined as number of tosses required to obtain 5th head when an unbiased coin is tossed repeatedly under identical conditions. Find a) probability generating function of Y.

b) E(Y) and V(Y).

Let $X_1, X_2, X_3, \dots, X_{12}$ be mutually independent random variables having common probability mass function

 $p(x) = (1/4)^{x} (3/4)^{1-x}; x = 0, 1.$ = 0; otherwise

Find

- a) probability generating function of X i.
- b) probability generating function of Y.

c) E(Y) and V(Y)

where $Y = X_1 + X_2 + X_3 + X_{12}$

- Three fair six sided dice are tossed simultaneously. Let X denote the sum of the numbers observed on the uppermost faces of the 3 dice. Find
 - a) probability generating function of X.
 - b) P(X = 17).
 - c) P(X = 6).
- 6) Four tickets are drawn at random with replacement from a bag containing ten tickets numbered 1, 2,..., 10. Let X denote the sum of numbers on the tickets drawn. Find:
 - a) probability generating function of X
 - b) P(X = 38).
- 7) In a lottery, tickets are numbered from 000000 to 999999 and they are equally probable. What is the probability that the sum of the digits of a chosen number will be 21?

8) The joint p.d.f. of X and Y is given below

	0	1	2
0	1/16	1/8	1/16
1	1/8	1/4	1/8
2	1/16	1/8	1/16

Find

- a) probability generating function of X.
- b) probability generating function of Y.
- c) probability generating function and probability distribution of Z = X + Y.

T.Y.B.Sc. Practical No.6.1.3 STOCHASTIC PROCESSES

The time between arrivals at L and J Restaurant is exponential with mean 5 minutes. The restaurant The nine business at 11.00 a.m. Determine the following opens for business at 11.00 a.m. Determine the following opens for business.

a) The probability of having 10 arrivals in restaurant at 11.12 a.m. given that there were 8 at 11.05

a.m.
b) The probability that a new customer will arrive between 11.28 and 11.33 a.m. given that the is customer arrived at 11.25 a.m.

customer and control of the process with initially 2 many.

c) The probability of birth process with initially 2 members, experiences an average birth rate of 3 A Yule type Obtain members per hour. Obtain

memoris possibility of having 4 or more members in the population after 1 hour 15 minutes, a) probability of having 4 or more members in the population after 1 hour 15 minutes,

a) probability

a) probability

b) expected size and standard deviation of population at 2 hours and 30 minutes.

b) expected an item occurs according to a Poisson distribution with mean 2 per day. The maximum pentalio 10 items, which occurs on each Monday immediately after a new order is received. The order size depends on number of units left at the end of the week on Saturday (business is closed on Sunday). Determine the following

a) The average weekly size of order.

b) The probability of incurring shortage when the business opens on Friday morning.

- A population consists of elements which can split or die. The rate of splitting in a two is 3 per unit A population of dying is 2 per unit time. If there is no interaction among the elements, obtain the mean and standard deviation of population after 2 time units if initial population size is 100.
- An office opens at 11 a.m. daily. There are 2 particular employees who come before 11 a.m. and other employees arrive at the rate of 2 employees per minute after 11 a.m. Obtain the probability 5) that there are

b) 10 employees at 11.05 a.m. Also obtain expected number of employees at 11.02 a.m.

A machine shop has just stocked 10 spare parts for the repair of a machine. Stock replenishment of size 10 pieces occurs every 7 days. The time between breakdowns is exponential with mean 1 day. Determine the probability that the machine will remain broken for 2 days because of the unavailability of parts.

T.Y.B.Sc. Practical No.6.1.4

QUEUEING THEORY-I

- 1) Customers arrive at the first class ticket counter of a railway station at a rate of 12 per hour. There is one clerk serving the customers at a rate of 30 per hour. Obtain a) the probability that the system is empty. b) the probability that there are more than 2 customers in the system. c) the probability that there is no customer waiting to be served, d) the probability that a customer is being served and exponentially distributed.
- A tailor specializes in ladies' dresses. The number of customers approaching the tailor appears to be Poisson distributed with a mean of 6 customers per hour. The tailor attends the customers on a first-to attend one customer is 6 minutes. Obtain a) the average time that the tailor is free on a 10-hour working day. b) the expected number of customers in the tailor's shop, c) the expected number of customers waiting for tailor's services. d) the probability that an arriving customer will be waiting shop for more than 10 minutes. e) the probability that an arriving customer will be in the
- 3) At Dr. Prachi's clinic, on an average, 8 patients arrive per hour. Some patients require only the repeat prescription; others come for minor check-up, while some others require thorough inspection for the diagnosis. This takes Dr. Prachi six minutes per patient on an average. It can be assumed that distribution. Obtain a) the percent of times a patient can walk inside the doctor's cabin without having to wait. b) the average number of patients waiting for their turn. c) the average time that a patient spends in the clinic.
- 4) A post office has 3 windows providing the same service. It receives on an average 30 customers per hour. Arrivals are in Poisson type and service time is exponentially distributed. Each window serves on an average 12 customers per hour. Obtain a) the probability that an arriving customer will be served immediately. b) the probability that a customer will have to wait. c) the average number of customers in the system. d) the average total time that a customer spend in the post office.
- 5) A bakery is staffed by 2 clerks, each of them capable of handling an average of 30 customers an hour. Customers arrive at the bakery according to a Poisson process, at a mean rate of 50 per hour and the service time is exponentially distributed. Obtain a) the probability that there are more than 3 customers in the bakery at the same time. b) the probability that a given clerk is idle.
- 6) A to Z Groceries operates with three check-out counters and the operation is set up such that the customer will go to the first empty counter. Customers arrive at the store according to a Poisson distribution, with a mean of 10 per hour. The service time per customer is exponential with mean 12 minutes. Obtain a) the expected number of customers waiting to be served. b) the expected number of busy servers. c) probability that all the three counters are busy.
- On an average, 96 patients per 24-hour day require the service of an emergency clinic. Also, on an average, a patient requires 10 minutes of active attention. Assume that the facility can handle only one emergency at a time. Suppose that it costs the clinic Rs.100 per patient treated. Each minute of decrease in the average service time would cost Rs.10 per patient treated. How much would have to be budgeted by the clinic to decrease the average size of the queue from 4/3 patients to ½ patients?
- 8) A company currently has two tool cribs, each having a single clerk, in its manufacturing area. One tool crib handles only the tools for heavy machinery, while the second one handles all other tools. It is observed that for each tool crib the arrivals follow a Poisson distribution with a mean of 20 per hour and the service time distribution is exponential with a mean of 2 minutes.
 - The tool manager feels that if the tool cribs are combined in such a way that either clerk can handle any kind of tools as demand arises, it would reduce the waiting time in the system.

 Analyze the data and comment on the management's opinion.

T.Y.B.Sc. Practical No.6.1.5

QUEUEING THEORY-II

A small barber shop, operated by a single barber, has a room for at most 2 customers. Potential customers arrive in a Poisson fashion at an average rate of 3 per hour. Service time follows exponential distribution with mean 1/4 hour. Obtain

a) the average time spent in the shop by an entering customer.

b) the probability that an arriving customer can enter the system.

c) the probability that the barber is busy.

Customers arrive at a beauty parlour in a Poisson fashion at an average rate of 6 per hour. Any arriving customer is denied service if there are already 7 customers in all at the parlour. A single beautician attends to arriving customers on a FCFS basis at an average rate of 40 minutes per customer. The actual service time is exponential. Obtain

a) the percentage of time the beautician is busy.

b) the expected number of customers at the parlour.

c) the average time a customer will spend at the parlour.

d) the average number of customers who are denied entry per hour.

A service center is manned by 5 servers each of whom work independently at an average rate of 15 minutes per customer. The actual distribution of service time for each server is exponential. Customers arrive at the center in a Poisson fashion at an average rate of 10 per hour. Any arriving customer is forced to leave the system if all the servers are busy. Obtain

a) the steady state probability distribution of the number of customers in the system.

b) the probability that 2 or more servers are free.

c) the expected waiting time in the system for any arrival.

4) A self service car wash has four stalls in which customers can clean and wax heir automobiles and room to accommodate maximum of three additional cars when all stalls are full. Customers arrive at the car wash according to Poisson process at a mean rate of 15 per hour. If there is no room for them on the grounds of the car wash, arriving customers must go elsewhere. The time required to service a car appears to be exponentially distributed with a mean of 12 minutes. Obtain

a) the average number of cars at the car wash at any given time.

b) the expected rate at which cars are denied entrance to the car wash.

5) One repairman is attending 5 machines. The rate of breakdown per machine is one per hour while the service rate is ten per hour. Obtain

a) the probability that the repairman is idle.

b) the expected number of machines waiting for the service.

6) A company has five machines, each of which suffers breakdown at an average rate pf 0.5 per hour. There are two repairmen and average repair time taken by each of them is 30 minutes. Obtain a) the probability that all machines are in good condition.

b) the average time an out of order machine has to spend waiting for repairs to start.

7) A doctor's clinic can accommodate at most 10 patients (including the one being examined). Patients arrive at the clinic in a Poisson fashion at a rate of 5 per hour, the time taken by the doctor to examine one patient is exponentially distributed with a mean of 12 minutes per customer. Obtain a) the expected waiting time in the system of a customer.

b) the percentage of time the doctor is idle.

c) the expected number of customers per hour who are denied the service.

At a self-service facility arrivals occur according to Poisson fashion with mean 30 per hour. The distribution of service time is exponential with mean 5 minutes. Obtain

a) the expected number of customers in service at service facility.

b) the percentage of time the facility is not used by customers.

T.Y.B.Sc. Practical No.6.2.1

CHEBY CHEV'S INEQUALITY

The probability distribution of a random variable X is given by 1)

$$p(x) = 1/8$$
; $x = -1, 1$
= 6/8 : $x = 0$.

Obtain a bound to $P(|X| \ge 1)$. Compare it with the actual probability and comment on the same.

- 10% of a large lot of apples are bad. If N denotes the number of bad apples in a random sample of 2) size 20, obtain a bound to $P(0 \le X \le 4)$ and compare your answer with the actual probability.
- It is suspected that 40% persons of a large community have high blood pressure. What is the 3) Chebychev's lower bound to the probability that in a random sample of 100 persons from the community, 30% to 50% of them would test positive for high blood pressure? Obtain the above probability using Central limit theorem and Compare the two.
- The number of persons X_A and X_B , lined up respectively at railway booking windows A and B is 4) independent and Geometrically distributed with probability distribution given by

$$p(x) = q^{x}p$$
, $x = 0, 1, 2, 3, ...$

where p = 0.2 for window A and p = 0.5 for window B. An arrival joins the queue at any window at

- a) $P(|X_A E(X_A)| \ge 5)$.
- b) $P(|X_B E(X_B)| \ge 3)$.
- c) $P(|X E(X)| \ge 8)$ where $X = X_A + X_B$.
- The marks of any student in T.Y.B.Sc. 1as mean 60 and standard deviation of 4. Find bound to the
 - a) marks of a given student lies between 55 and 65.
 - b) the average marks of five given students lie between 55 and 65.
 - c) the absolute difference in marks of any two given students is greater than or equal to 6.
- (X_1, X_2, X_3) follows multinomial distribution with parameters (10, 0.2, 0.3, 0.4).
 - a) obtain Chebychev's bound to $P(0 \le X_1 \le 4)$, $P(3 \le X_1 + X_2 \le 7)$.
 - b) compare each of the above with actual probability.
- The chance of failure X of an electronic component has p.d.f. given by

$$f(x) = 504 \text{ x}^{5}(1-x)^{3}$$
; $0 \le x \le 1$
= 0 ; otherwise

- Obtain a bound to P(0.4 < X < 0.8).4
- The amount of toll X (in lakhs of rupees) collected per day at a booth is Uniformly distributed over the interval (20,40). Obtain Chebychev's bound to the probability that the average collection per day during the last one week lies between Rs. 26 lakhs and Rs. 34 lakhs.

T.Y.B.Sc. Practical No. 6.2.2

OFDER STATISTICS

- Given a random sample of size seven from a population of continuous type, obtain the probability 1)
 - a) smallest order stati ic exceeds the population median.
 - b) second order statistic is less than the population quartile Q₁.
 - b) second on the population quartiles Q_1 and Q_3 .
- Five observations are randomly made from a population with p.d.f.

2)
$$f(x) = 2(1-x)$$
; $0 < x < 1$
= 0; otherwise.

Find the probability that

- a) the sample median is more than 1/2.
- b) fourth order statistic is more than 1/4.
- A random sample of size six is drawn from a continuous Uniform distribution over the interval (0, 2). Obtain p.d.f. of the sample range and hence probability that the sample range is less than K
- The time taken per telephone call X in minutes follows an Exponential distribution with mean 5. Obtain the probability that the sample range exceeds 4 minutes, based on a random samples of size
- 5) The monthly income X in thousands of rupees of workers in an area has p.d.f. $f(x) = e^{-(x-\theta)}$; $x > \theta$, $\theta > 0$

$$f(x) = e^{-(x-\theta)}$$
; $x > \theta$, $\theta > 0$
= 0; otherwise

A random sample from the above population is as follows 12, 10, 15, 12, 11, 10, 11, 17, 9, 22. Obtain at unbiased estimate of θ based on the smallest order statistic and standard error of your

- 6) There is a bus service every θ minutes where θ is unknown. The time interval between any two consecutive services is Uniform continuous over the interval $(0, \theta)$. The time intervals between any two consecutive services are recently observed to be as follows: 5, 7, 6.5, 4, 4, 3, 6, 8, 5.5, 6.
 - a) an unbiased estimate of θ based on the order statistics (i) Y_5 (ii) Y_{10} .
 - b) standard errors of your estimates.
 - c) correlation coefficient between Y₅ and Y₁₀.
- 7) A random sample of size two is drawn from a population with p.d.f.

$$f(x) = 2x ; 0 < x < 1$$
= 0 ; otherwise

Find the probability that one of them is at least twice as large as the other.

- 8) A random sample of size three is drawn from an Exponential distribution with mean one. Obtain
 - b) mean and standard deviation of Y₁.

B.Sc. Practical No.6.2.3

BIVARIATE DISCRETE DISTRIBUTIONS

Let W be the weight of laundry soap in 1-kilogram box, $P(W \le 1) = 0.02$ and $P(W \ge 1.072) = 0.08$. Let w to so of soap light, good or heavy depending on whether W < 1, $1 \le W \le 1.072$. We call a box of soap light, good or heavy depending on whether W < 1, $1 \le W \le 1.072$. We can a solution with the sympletrian of good boxes of W > 1.072, respectively. In a random sample of E = 50 boxes, let X denote the number of light E = 1.072, respectively. In a random sample of E = 50 boxes, let X denote the number of light boxes and Y the number of good boxes. a) what is the joint p.d.i. of X and Y? b) what are the parameters of the distribution? c) given that X = 3, how is Y distributed conditionally? d) determine E(Y / X = 3). e) find p, the correlation coefficient between X and Y. It is observed that the percentage of students securing first class, second class, pass class and failure at a certain examination are 40, 20, 30 and 10 respectively out of 100 students who have appeared for this examination. Let X_1 , X_2 and X_3 be the number of students who have secured 1^{st} , 2^{nd} and pass class respectively. State a) the joint distribution of X_1 . X_2 and X_3 . b) marginal distribution of X1, X2 and X3. c) joint marginal distribution of X_1 and X_2 . Hence find correlation between X_1 and X_2 . d) conditional distribution of X1 and X2 for given X3. e) conditional distribution of X_1 for given X_2 and X_3 hence find $E(X_1/X_2=10 \text{ and } X_3=15)$. The M.G.F. of (X, Y) is given by $(0.2e^{11} + 0.5e^{12} + 0.3)$ Write the probability distribution of a) (X, Y). b) Y given X = 2. c) X given Y = 2. Also find E(Y / X = 2) and E(X / Y = 2). One of the numbers 1, 2,..., 6 is to be chosen by casting an unbiased die. Let this random experiment be repeated five independent times. Let the random variable X1 denote the number of terminations in the set $\{x ; x = 1, 2, 3\}$ and let the random variable X_2 denote the number of terminations in the set $\{x; x = 4, 5\}$. Compute P($X_1 = 2, X_2 = 1$).

terminations in the set
$$\{x; x = 4, 5\}$$
. Compute $Y(X) = 4$. Four independent observations are made on a random variable X having p.d.f.
$$f(x) = 1 - |x|; -1 < x < 1$$

$$= 0$$
otherwise

= 0; otherwise Suppose that the interval -1 to 1 is divided into 4 distinct classes of the same length. Find the probability that one observation falls in each of the first two classes, two in the third and none in the last class.

T.Y B.Sc. Practical No.6.2.4

BIVARIATE CONTINUOUS DISTRIBUTIONS

1) Let X and Y have a Bivariate Normal Distribution with parameters $\mu_1 = 3$, $\mu_2 = 0$, $\sigma_1^2 = 16$, $\sigma_2^2 = 25$, $\rho = 0.6$

Determine the following probabilities.

a)
$$P(2 \le Y \le 7)$$

 $y \ge 1(2 \le Y \le 7 / X = 7)$

c)
$$P(-3 \le X \le 3)$$

d)
$$P(-3 < X < 3 / Y = -3)$$

2) Let X and Y have a Bivariate Normal Distribution with parameters.

$$\mu_1 = 1$$
, $\mu_2 = 2$, $\sigma_1^2 = 16$, $\sigma_2^2 = 25$, $\rho = 12 \cdot 13$

Determine the following probabilities.

a)
$$P(X > 2)$$

b)
$$P(X > 2 / Y = 2)$$

3) A Bivariate Normal Distribution has the p.d.f.

$$f(x,y) = C \exp \left[-x^2 + xy - y^2\right]; -\infty < x,y < \infty$$

Find the constant C and the parameters of the distribution. Also find the conditional p.d.f. of Y given X = 2.

4) Find the constant k and the parameters of the following Bivariate Normal Distribution.

$$f(x,y) = k \exp \left\{ -\frac{8}{27} [(x-7)^2 - 2(x-7)(y-5) + 4(y+5)^2] \right\}$$

5) A Bivariate Normal Distribution has the p.d.f.

$$f(x,y) = k \exp \left\{ -\frac{1}{102} \left[(x+2)^2 - 2.8(x+2)(y-1) + 4(y-1)^2 \right] \right\}$$

Find

- a) μ_1 , μ_2 , σ_1 , σ_2 , ρ .
- b) conditional mean of Y given X = x.
- c) V(Y/x)
- 6) The marks X and Y scored by candidates in an examination in two subjects are known to follow a Bivariate Normal Distribution. The mean of X is 52 and its s.d. is 15 while Y has mean 48 and s.d. 15. Also ρ = 0.6. Write down the joint p.d.f. of X and Y. If 100 marks in the aggregate are needed to pass in the examination, find the proportion of candidates who pass in the examination. Also find P(|X-Y| < 10).</p>

Y.B.Sc. Practical No.6.2.5

SIGNIFICANCE OF CORRELATION COEFFICIENT

A random sample of 30 pairs of observation from a normal population gave a correlation A random sample of 0.7. Is it likely that the variables in the population are uncorrelated? what is the least value of 'r' in a random sample of 27 pairs that is significant a) at 0.05 leve:

b) at 0.01 level.

The height (in cms) and the number of leaves of 15 sunflower plants was obtained and recorded as follows. Calculate correlation coefficient and test its significance.

Ht (in cms)	20	27	29	24	30	37	23	24	28	15	25	31 25	37	21
No.of leaves	16	19	18	20	25	30	16	29	26	12	18	23 24	28	19

A correlation coefficient of 0.72 was obtained from a sample of 29 pairs of observations.

a) Can the sample be regarded as drawn from a bivariate normal population with correlation coefficient of 0.8?

b) Obtain 95% confidence interval for the population correlation coefficient p in the light of the information provided by the sample.

In an air pollution study a researcher obtained the following results with regard to the number of lichen species (X) and the respective distance from each of the 25 chemical factories (Y in kms). $\Sigma X = 1165$, $\Sigma Y = 185.20$, $\Sigma X^2 = 56947$, $\Sigma Y^2 = 1437.24$, $\Sigma \Sigma XY = 9024.40$. Test the hypothesis that the corresponding correlation value for the population is 0.8.

The correlation coefficient between body length and body weight of Anabas scandens (tresh water fish) for two samples of sizes 40 and 50 was 0.5 and 0.55 respectively. Test whether the correlation coefficients are significantly different.

T.Y.B.Sc. Practical No.6.2.6

MULTIPLE AND PARTIAL CORRELATION AND REGRESSION

following table gives the weights X1 (in pounds), heights X2 (in inches) and the ages of 12

• • •	The "
1)	boys.

Weight X1	64	71	53	67	55	58	7.7	57	56	51	76	78
Height X2		59	49	62	51	50	55	48	52	42	61	57
Age X ₃	8	10	6	11.	8	7	10	9	10	6	12	9

a) Find the least square regression of X1 on X2 and X3.

b) Estimate the weight of a boy who is 9 years old and 54 inches tall.

b) Estimate and a Pizza shack collected the following data for the past 12 months on the total The manager of a Pizza shack collected the following data for the past 12 months on the total the manager of a pizza shack collected the following data for the past 12 months on the total The manager X_1 (in 000's of rupees), number of advertisements appearing every month (X_2) and monthly sales X_1 (in 000's of rupees) monthly cost of advertising (in 00's of rupees).

$$\sum X_{1} = 432.4 \qquad \sum X_{2} = 110 \qquad \sum X_{3} = 133.1$$

$$\sum X_{1}^{2} = 16034 \qquad \sum X_{2}^{2} = 1072 \qquad \sum X_{3}^{2} = 1513.27$$

$$\sum \sum X_{1} X_{2} = 40963 \qquad \sum \sum X_{1} X_{3} = 4902.23 \qquad \sum \sum X_{2} X_{3} = 1263.5$$

a) Find the plane of regression of X1 on X2 and X3.

b) Estimate the total monthly sales of pizza if the cost of advertisement is Rs.15,200 and the number of advertisement monthly is 16.

3) An instructor wished to determine the relationship of marks on a final examination to those on the two unit tests conducted during the year. Calling X₁, X₂ and X₃ as the marks of a student on the final exam, first unit test and second unit test respectively, he obtained the following computations for a total of 120 students.

$$\overline{X}_1 = 73$$
, $\overline{X}_2 = 6.8$, $\overline{X}_3 = 7.0$, $\sigma_1 = 9.0$, $\sigma_2 = 1.0$, $\sigma_3 = 0.8$
 $\sigma_{12} = 0.65$, $\sigma_{13} = 0.60$, $\sigma_{23} = 0.70$

a) Find the least square equation of X_1 on X_2 and X_3 .

b) Estimate the final marks of two students who secured respectively (9, 7) and (4, 8) in the two

c) Obtain multiple correlation coefficient R $_{1,23}$ and test its significance.

d) Obtain partial correlation coefficient r_{12.3} and test its significance.

In a tri-variate distribution, it is given that

$$\sigma_1 = 3$$
; $\sigma_2 = 4$; $\sigma_3 = 5$; $r_{23} = 0.9$; $r_{13} = 0.6$; $r_{12} = 0.65$

a) Obtain regression line of X_1 on X_2 and X_3 .

b) Find Partial correlation coefficients r 12.3, r 13.2 and r 23.1

c) Find Multiple correlation coefficients R 1.23, R 2.12 and R 3.12

d) Calculate $\sigma^2_{1,23}$, $\sigma^2_{2,13}$, $\sigma^2_{3,12}$

(Note the variates are measured from their respective means.)

Y.B.Sc. Practical No. 7. 1. 1 B.SC. PORTALIMUM LIKELIHOOD, METHOD OF MOMENTS

NETHOD OF MODIFIED MINIMUM CHI-SQUADE AND METHOD OF MODIFIED MINIMUM CHI-SQUARE

Let probability density function (p.d.f.) of a random variable (r.v.) X be f(x, \theta), and a random and of observations from a population with this p.d.f. is given. Estimate the unknown Let probability density function (plant) of moments and a random variable (r.v.) X be f(x, 0), and a random sample of observations from a population with this p.d.f. is given. Estimate the unknown parameter (i) method of moments

(i) method of maximum likelihood (i) method of maximum (ii) method of maximum error of the m.l.e, for each of the following examples (a),(b),(c):

and estimate standard error of the m.l.e, for each of the following examples (a),(b),(c):

otherwise. and estimate $\theta = \theta = \theta = 0$; x > 0, $\theta > 0$ a) $f(x, \theta) = 0$; otherwise.

b) $f(x, \theta) = \theta x$; $0 < x < 1, \theta > 0$.

 $\begin{array}{c} 10^{3} = 0 \\ \text{Observed sample values : 0.41, 0.57, 0.75, 0.78, 0.65, 0.82, 0.48, 0.39, 0.65, 0.51,0.46, and 0.83.} \\ x = -x/\theta + y > 0. \theta > 0. \end{array}$

 $c) f(x, \theta) = \frac{x}{\theta^2} e^{-x/\theta} ; x > 0, \theta > 0.$

Observed Sample values: 5.4, 7.1, 5.6, 6.2, 6.4, 4.9, 5.8, 6.3, 5.5 and 4.8

Observed Sample of Sample of Sample of Observations from a population with Given the p.d.f. $f(x, \theta)$ of a r.v. X, and a random sample of Observations from a population with (i) method of moments

(i) method of maximum likelihood

for each of the following examples (a),(b),(c):-

a)
$$f(x, \theta) = \frac{1}{\theta}$$
; $0 < x < \theta$
= 0 : otherwise

= 0; otherwise.

Observed sample values are: 1.3, 0.6, 1.7, 2.2, 0.3 and 1.1.

b) $f(x, \theta) = e^{-(x-\theta)}$; $\theta < x < \infty, -\infty < \theta < \infty$

; otherwise.

Observed sample values are: 1.3, 0.7, 3.6, 5.4, 7.4, 2.5, 3.2, 4.1, 2.5 and 2.6.

$$f(x, \theta) = \frac{1}{2} e^{-|x-\theta|}; -\infty < x < \infty, -\infty < \theta < \omega$$

$$= 0$$
; otherwise.

Observed sample values are: 1.9, -1.8, 2.9, -2.2, 1.5, 2.2, -1.4, -1.5, 1.6 and 1.7.

The following table gives probabilities and observed frequencies in four classes AB, Ab, aB and ab in a genetic experiment

oxyouthent.					
CLASS	AB	Ab	аВ	ab	Total
Probability			a riggin i T		
1 Toolaumty	$2+\theta$	<u>1 – 0</u>	<u>1 – 0</u>	θ	1
Observed	4	4	4	4	
Frequency	108	30	25	7	170
					1

Where $0 < \theta < 1$. Estimate the unknown parameter θ by h) method of modified minimum chi-square

method of maximum likelihood and obtain its standard error.

The p.m.f. of a discrete random variable X is given by
$$p(x) \qquad \frac{2pq}{1-q^2} \qquad \frac{p^2}{1-q}$$

where q = 1 - p, 0 . A sample of 30 observations gave the frequencies for the two cases aswhere q 19 respectively. Estimate p using 11 and 19 respectively minimum p using a) method of modified minimum chi-square

a) method of maximum likelihood

c) method of moments. c) method of monators and their respective probabilities (as per a genetic model) are classified 1000 observations and color-blindness as follows: 1000 observation and color-blindness as follows:

anoies.	MALE	FEMALE
Obs. frequencies	440	515
NORMAL NORMAL	40	5
TOTIK DILLA		

Probabilities	MALE	FEMALE
NORMAL	(p/2)	$(p^2/2) + pq$
COLOR-BLIND	(q/2)	$(q^2/2)$

Given p + q = 1 and 0 , estimate parameter p by method of maximum likelihood. Also findexpected frequencies.

A random sample of 10 observations from a population with p.d.f.

$$f(x, \mu) = 2\mu x e^{-\mu x^2}$$
; $x > 0, \mu > 0$
= 0; otherwise

is given as follows: 1.2, 3.8, 6.8, 2.87, 0.75, 3.19, 0.72, 1.05, 5.4 and 7.82. Estimate unknown

parameter μ by

a) method of maximum likelihood and find its standard error

b) method of moments.

LY, B.Sc. Practical No. 7, 1, 2

METHOD OF MAXIMUM LIKELIHOOD AND METHOD OF MOMENTS

1) Let a random sample of 5 observations 8.1, 0.2, 1.6, 5.2, 2.1 be taken from a population with p.d.f. $\alpha \leq x \leq \beta$, $-\infty \leq \alpha$, $\beta \leq y$ $f(x, a, b) = \frac{1}{\beta - \alpha} : \alpha < x < \beta, -\infty < \alpha, \beta < \alpha$

Estimate parameters & and B by

Estimate parameters and b) method of maximum likelihood.

a) method of months and the months are model.

Let sample observations 2.3, 1.2, 0.9, 3.2, 1.6, 2.0, 2.9 and 3.4 be taken from a population with p.d.f $f(x, a, b) = \frac{1}{b} e^{-(x-a)/b} \; ; \; a < x < \omega \; , \; -\infty < a < \omega \; , \; b > 0$

Estimate parameters a and b by

a) method of moments and b) method of maximum likelihood.

a) method of the sample of size 10 was taken from a normally distributed population having mean μ and variance A sample of size $\chi \sim N(\mu, \sigma^2)$. The observations are as follows: 6.18, 5.02, 5.55, 6.32, 3.57, 4.82, 4.95, 5.80, 5.22 and 4.99. Find

a) maximum likelihood estimate (m.l.e) of (i) μ when $\sigma^2 = 0.4$ (ii) σ^2 when $\mu = 5.25$. b) m.l.e's of μ and σ^2 and P(X > 4.25).

The following data is drawn from a Negative binomial distribution with parameters k and p, 3 4

5 Total 4 1

Frequency: 214 123 45 13 400. Given $k \in \mathbb{N}$, 0 .Estimate parameters k and p by method of moments.

A random variable X takes values 0, 1, 2, 3, 4 and 5 with probabilities:

$$p(x) = (1 + pm) e^{-m} when x = 0$$

$$= m(1 - p) e^{-m} when x = 1$$

$$= \frac{e^{-m} m^{-x}}{x!} when x = 2, 3, 4, 5. Giver, m > 0 and 0$$

The observed frequencies are 37, 19, 22, 15, 6 and 1 respectively. Find maximum likelihood estimates of unknown parameters p and m.

BAYES' ESTIMATION

- The proportion θ of defective items in a large shipment is unknown and the prior distribution of θ is The proportion $\alpha = 5$ and $\beta = 10$. Suppose that 20 items are selected at random from the Bela with parameters with parameters one item is found to be defective. If the squared error loss function is shipment is the Bayes' estimate of θ? supplied, what is the Bayes' estimate of θ ?
- the daily number of incoming calls at a small city telephone exchange has a Poisson distribution.

 The daily number of incoming calls at a small city telephone exchange has a Poisson distribution. The daily number θ , however, varies from day to day. This variation is represented by a with mean $\alpha\beta=100$ and variance 200. Five days one at the second stribution is represented by a with mean $\alpha\beta$ =100 and variance 200. Five days are selected at random and the gamma density and the gamma density incoming calls found was 90, 95, 108,100 and 106. Find Bayes' estimate of the average number of daily incoming calls θ , using squared error loss function.
- white pine is one of the best known species of pines in the Northern United States and Canada. White pine is susceptible to a blister rust, which develops canker in the bark. These cankers swell, White place is stated in a forest. The number of discussion in the bark. These cankers swell, resulting in the death of twigs and small trees. A forester wishes to estimate the average number of diseased pine trees per acre in a forest. The number of diseased trees can be modeled by a Poisson distribution with parameter θ . Since θ changes from area to area, the forester believes that θ has an exponential distribution with mean $1/\alpha = 5$. Find Bayes' estimate of the mean number of diseased pine trees per acre, if the forester takes a random sample of 50 and
 - $\sum_{i=1}^{50} x_i = 200$. (Use squared error loss function).
- The distribution of lifetimes of bulbs made by a certain manufacturer is exponential with mean 1/0. Suppose θ has a priori exponential distribution with mean 1. Using squared error loss function, find Bayes' estimate of θ if the total of 25 observations is 29.
- The time (in minutes) required to serve a customer at a certain facility has an exponential distribution, for which the value of the parameter θ is unknown. The prior distribution of θ is a gamma distribution for which the mean is 0.2 and the variance is 1. The average time required to serve a random sample of 20 customers is observed to be 3.8 minutes. If squared error loss function is used, what is the Bayes' estimate of θ ?
- 6) The brain weight (in grams) of adults is assumed to have a normal distribution, that is
 - $N(\theta, \sigma^2 = 6)$. In order to estimate θ , a random sample of 150 gave $\sum_{i=1}^{150} x_i = 241380$ grams. The
 - experimenter believes that θ itself has a normal distribution N(1800, 9). Find Bayes' estimate of θ using squared error loss function. What is the Bayes' estimate if absolute error loss function is
- Given that 1, 0, 1, 1, 0, 0, 1, 0, 0, 1 is a random sample from a Bernoulli distribution with parameter θ where θ has prior uniform distribution on [0, 1], obtain Bayes' estimate of θ using squared error loss function. Also find the Bayes' risk.
- X = 6 is a single observation from a negative binomial, that is, NB(10, θ) distribution. If the prior distribution are less function. distribution of θ is Beta B(3, 5) obtain Bayes' estimate of θ using quadratic error loss function.

INTERVAL ESTIMATION

The results of 64 determination of strength (x) of a magnetic field steld a

ands with 1) $S(x_1 - \hat{x})^2 = 0.0063$. Obtain 90% confidence interval for mean μ assuming Normal distribution of X having an unknown variance.

Normal distribution with average yield and yield of strawberry plant tollows Normal distribution with average yield and Yield of strawberry plants in an uniformity trial are given below. Plani 10 no.

239 176 235 217 234 Yield 216 318 190 Compute 95% confidence limits for average yield μ when 101 225

a) σ^2 is known as 1600.

b) σ ² is unknown.

The following are the data on time X (in seconds) required to format 1.2 mb floppy diskettes on PC 386.

Floppy no. 1 2 3 6 8 100 90 97 101 93 95 Time 90 99

Assuming Normal distribution of X compute 99% confidence interval for variance of time for formatting floppy diskettes when

a) mean time is known as 90 seconds.

b) mean time is unknown.

4) A random sample of size 9 from $N(\mu, \sigma^2)$ population gives sample mean $\bar{x} = 68$ and $\sum (x_1 - \overline{x})^2 = 30$. Find 90% confidence interval for σ 2 if

a) μ is unknown.

b) u is known as 72.

5) A nutrition is examined the basal metabolism of 30 volunteers in two groups of m =16 and n = 14. He reported $\bar{x} = 36$ cal/sqm/hr, $\bar{y} = 32$ cal/sqm/hr. $\sum (x_i - \bar{x})^2 =$

72.86, $\sum (y_j - \overline{y})^2 = 14.46$. Obtain 95% confidence interval for $\frac{\sigma_x^2}{\sigma_y^2}$ assuming

Normal distribution of the basal metabolism in two groups with unknown means.

The oil contents of two brands of soap measured (in mg per gm) were found to be as follows

X 22 24 25 26 20 23 24 21 20 26 22 25 28 29 30 27

Assuming that oil contents X, Y are following N(μ_1 , σ_1^2) and N(μ_2 , σ_2^2) find 99%

confidence interval for $\frac{\sigma_1^2}{\sigma_2^2}$.

Let X denote the permeability of toroids in upper deck of furnace and Y that in the lower deck. An experiment was performed and ε bservations on permeability of toroids were obtained. Permeability of toroids is as follows toroids were obtained. Permeability of 1964.2 1848.0 2129.6 2218.4 2236.8 X 1919.5 1916.7 1891.6 1964.2 1848.0 2129.6 2218.4 2236.8 Y 2143.4 2055.0 1922.0 2068.1 1911.0 19"4.4 2055.1 2154.8 Compute 95% confidence interval for difference between averages of two populations (assume normality.

populations (assumptions) $x \ge 0$, $\theta > 0$. Using m.l.e of θ and asymptotic distribution of Let $f(x, \theta) = \theta e^{-\theta x}$; $x \ge 0$, $\theta > 0$. Using m.l.e of θ and asymptotic distribution of m.l.e, obtain 99% confidence interval for θ if sample mean of 43 observations is

13.73.

T.Y.B.Sc. Practical No.7.1.5

LINEAR MODELS

A soft drink bottler is analyzing the vending machine service routes in his distribution system. The A soft drink countries that the two important variables affecting the engineer responsively) are the number of cases of product stocked(x₁), and the distance walked by the delivery time(x₂). The engineer has collected 10 observations on delivery time(x₂). delivery time(y) and the distance walked by the outer driver(x 2). The engineer has collected 10 observations on delivery time which are shown

route							Tallian 19w		are siton	VII
below. obs** no. delivery time:y no.of cases:X 1 distance (ft):X 2	560	$ \begin{array}{c} 2\\11.50\\3\\220\\\text{del } \mathbf{v} = 6 \end{array} $	$\frac{3}{12.03}$ $\frac{3}{340}$	4 14.88 4 80 X 1 + B a	5 13.75 6 150	6 18.11 7 330	7 8.00 2 110	8 17.83 7 210	9 79.24 30 1460	10 21.50 5 605
C'A A HITCAL SWA				- 1 P 2	A 7 1 C.	IO IDE A	PITTION	· · · · · · · · · · · · · · · · · · ·		

distance (1).32 (distance (1).32) (110 210 1460) (distance (1).32) (110 210 1460) (a) Fit a linear statistical model $y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + e$, to the delivery time data, such that $y_i \sim N(E(y_i), \sigma^2)$ and $e_i \sim N(0, \sigma^2)$ for i = 1, 2, ..., 10. $\beta_0, \beta_1, \beta_2$ are unknown parameters. b) Find best linear unbiased estimate of β_0 , β_1 , β_2 .

c) Also find an estimate of σ^2 and of $cov(\hat{\beta}_1, \hat{\beta}_2)$.

d) Obtain 95% confidence interval for β_1 when (i) $\sigma^2 = 8.8$ (ii) σ^2 is unknown.

An experiment was conducted to determine the influence of acid bath temperature(x), on an appropriate measure of the whiteness of rayon (y). X is the controlled variable and Y is the

dependent variable.

White	ness of	Temperature	
y_1	y_2	у ₃	X
76.5	80.2	76.0	30
83.5	89.1	79.9	35
84.2	77.2	89.5	40
85.7	85.1	99.5	25
97.5	71.5	89.4	45
103.2	84.5	108.7	50
115.2	77.5	111.5	35
108.1	79.2	102.3	65

If 7_i is the average-whiteness at acid bath temperature (x_i), and if we assume that

$$E(\bar{y}_i) = \beta_0 + \beta_1 x_i \text{ for } i = 1, 2, ..., 8.$$

where β_0 and β_1 are unknown parameters. Find

a) the least square estimate of β_0 and β_1 .

b) 99% contidence interval for β 0.

c) 98% confidence interval for σ^2 .

The following gives a property evaluation data, where y: sale price of the house (in Rs 1900):

x₁: living space(square feet); x₂: age of houses (in years).

V	25.0	00.	1001),	x 2 . age	ot nous	es (in ve	ears)			,		
,	~~./	27.3	770	250	0.5	-	,					
x 1	993	1500		25.5	29.9	29.9	30.9	28.9	35 0	31 6	21.0	
٧.		1000	11/5	1232	1121	000	1210	1.70-	.,,,	31.5	31.0	30.9
A 2	4]	62	40	-202	1141	200	1240	1501	1225	1552	075	1101
a) Fit the	· t	02	4(;	54	42	56	5.1	2.2	20	31.5 1552	213	1121
7 - 11 [1]	unear	Statistic:	l madel		.~	20	31	32	32	1552 30	30	2.3

a) Fit the linear statistical model $y = \beta_0 + \beta_1(\bar{x}_1 - \bar{x}_1) + \beta_2(\bar{x}_2 - \bar{x}_2) + e$

where $y_i \sim N(E(y_i), \sigma^2)$ and $e_i \sim N(0, \sigma^2)$ for i = 1, 2, ..., 12. b) Find best linear unbiased estimate of β_0 . β_1 , β_2 .

90btain the best linear unbiased estimate of $\beta_0 + 2\beta_1 - \beta_2$ and its standard error. 4) Obtain 95% confidence interval for $\beta_0 + 2\beta_1 - \beta_2$.

e) Test the hypothesis H_0 : $\beta_0 + 2 \beta_1 - \beta_2 = 33$.

18 Sc. Practical No. 7, 2, 1 TESTING OF HYPOTHESES-1

The length of a tube has exponential p.d.f. with mean θ . In order to test H_0 : $\theta \neq 10$ against the length of a tube has exponential region of the form $C = \{X: X \leq c_1 \text{ or } X \geq c_2\}$ where The length of a tube has copour and probability and the size of the test is 0.05. What is the point θ and power at $\theta = 2$. 8. 15. 20 The fermion consider the critical value of which an unknown of the last? Find power at $\theta = 2$, 8, 15, 20.

and a such time test? Find power at $\theta = 2, 8, 15, 20$, function of the test? Find power at $\theta = 2, 8, 15, 20$. and of the test rand power of the test rand Induce of are white. To test the hypothesis and an arm contains 0 markets at random from the He $0 \le 3$ against $0 \le 3$ and $0 \le 3$ against $0 \le 3$ against $0 \le 3$ and $0 \le 3$ against $0 \le 3$ against $0 \le 3$ against $0 \le 3$ and $0 \le 3$ against $0 \le 3$ against $0 \le 3$ and $0 \le 3$ against $0 \le 3$ against $0 \le 3$ and $0 \le 3$ against $0 \le 3$ against $0 \le 3$ and $0 \le 3$ against $0 \le 3$ against $0 \le 3$ and $0 \le 3$ against $0 \le 3$ against $0 \le 3$ and $0 \le 3$ against $0 \le 3$ against $0 \le 3$ and $0 \le 3$ against $0 \le 3$ against $0 \le 3$ and $0 \le 3$ against $0 \le 3$ against $0 \le 3$ against $0 \le 3$ and $0 \le 3$ against $0 \le 3$ Here 3 against rules and reject H_0 if one of them is white. Find the level of significance and without replacement at $\theta = 5$, 10, 18.

power of the test at o cell as of type A or type B is long and expensive, but recent series of the identification of a cell as of type A or type B is long and expensive, but recent series of the identification of a cell as of type A or type B is long and expensive, but recent series of the identification of a cell as of type A or type B is long and expensive, but recent series of the identification of a cell as of type A or type B is long and expensive, but recent series of the identification of a cell as of type A or type B is long and expensive, but recent series of the identification of a cell as of type A or type B is long and expensive, but recent series of the identification of a cell as of type A or type B is long and expensive, but recent series of the identification of a cell as of type A or type B is long and expensive, but recent series of the identification of a cell as of type A or type B is long and expensive, but recent series of the identification of a cell as of type A or type B is long and expensive, but recent series of the identification of a cell as of type A or type B is long and expensive, but recent series of the identification of a cell as of type A or type B is long and expensive, but recent series of the identification of a cell as of type B is long and expensive and type B is long and expensive at the identification of the identificatio power of the test at $\theta = 5$, 10, 18. The identification of a contact that the quantity (in certain units) of an easily quantifiable chemical experiments have shown that the quantity (in $\mu = 10$, $\sigma^2 = 1$) random variable of the cell is well described by a N($\mu = 10$, $\sigma^2 = 1$) random variable. experiments have shown that the sho contained in the central variable for type B. In future it is decided to classify a cell as type A if $N\mu = 15$, $\sigma^2 = 4$) random variable for type B. In future it is decided to classify a cell as type A if $N\mu = 15$, $\sigma = 4$) random values of chemical and as type B otherwise. Obtain suitable measures of it contains not more than 12 units of chemical and as type B otherwise. Obtain suitable measures of the contains to misclassify. Where should the point of division be drawn if it is a solution of the contains to misclassify. it contains not more should the point of division be drawn if it is required to set the fine liabilities to misclassifying a type A cell equal to 0.01? What then is the misclassifying a type A cell equal to 0.01? rice liabilities to misclassifying a type A cell equal to 0.01? What then is the probability of other kind probability of misclassifying?

A random sample of size 2 is drawn from exponential distribution with mean θ to test H₀: $\theta = 2$ Arandom samp. The critical region is $C = \{ (X_1, X_2) / 0.5 \le X_1 + X_2 \le \infty \}$. Find the level of against $H_1: \theta = 3$. The critical region is $C = \{ (X_1, X_2) / 0.5 \le X_1 + X_2 \le \infty \}$. Find the level of

significance and power of the test.

Arandom variable X has the following probability density function

$$\frac{\text{dom variable } X}{f(x, \theta)} = \theta x^{\theta - 1}; \quad 0 < x < 1$$

$$= 0 \qquad ; \quad \text{otherwis}$$

To test H_0 : $\theta = 1$ against H_1 : $\theta > 1$ the critical region is $C = \{ (X_1, X_2) / 0.75 \le X_1 X_2 \}$. Find the size of the test and values of power function at $\theta = 4$, 6, 10, 15. Sketch the power curve.

An item is packed in lots of 100 each. Let X denote the number of defectives in a packet of 100 and is a Poisson variable with mean m. To test H_0 : m = 0.5 against H_1 : m < 0.5 it is decided to select 12 packets and inspect the items in them. If they contain less than two defectives, it is proposed to reject H₀. Find the level of significance of the test. Obtain the K(m) power function of the test Sketch the power curve, considering m = 0.3, 0.25, 0.16, 0.08, 0.02.

7) Consider a random variable following a Normal distribution with mean θ and variance 4. The simple hypothesis $H_0: \theta = -1$ is rejected and the alternative $H_1: \theta = 1$ is accepted if

 $X_1 + 2X_2 + 3X_3 + 4X_4 + \dots + 10X_{10} > 0$. Find the power and size of the test.

1) The proportion p of swimmers in a large population is unknown and it is desired to test $H_0: p=0.5$ against $H_1: p = 0.75$. A random sample of 5 people is drawn from the population. Let Y = number of swimmers in the sample. The following critical regions are proposed.

 $\emptyset C_1 = \{ Y \le 1 \}, \text{ ii) } C_2 = \{ Y \ge 4 \}, \text{ iii) } C_3 = \{ Y = 1,5 \}.$

In each case obtain probability of type I and type II errors and hence suggest the best critical region from the given three.

B.Sc. Practical No.7.2.2 TESTING OF HYPOTHESES-II

Suppose X 1, X 2, X 3, X 4, X 5 is a random sample from Bernoulli distribution with parameter p. Find suppose X 1, X 2, X 3, X 4, X 5 is a random sample from Bernoulli distribution with parameter p. Find suppose X 1, X 2, X 3, X 4, X 5 is a random sample from Bernoulli distribution with parameter p. Find suppose X 1, X 2, X 3, X 4, X 5 is a random sample from Bernoulli distribution with parameter p. Find suppose X 1, X 2, X 3, X 4, X 5 is a random sample from Bernoulli distribution with parameter p. Find suppose X 1, X 2, X 3, X 4, X 5 is a random sample from Bernoulli distribution with parameter p. Find suppose X 1, X 2, X 3, X 4, X 5 is a random sample from Bernoulli distribution with parameter p. Find suppose X 1, X 2, X 3, X 4, X 5 is a random sample from Bernoulli distribution with parameter p. Find suppose X 1, X 2, X 3, X 4, X 5 is a random sample from Bernoulli distribution with parameter p. Find suppose X 1, X 2, X 3, X 4, X 5 is a random sample from Bernoulli distribution with parameter p. Find suppose X 1, X 2, X 3, X 4, X 5 is a random sample from Bernoulli distribution with parameter p. Find suppose X 1, X 2, X 3, X 4, X 5 is a random sample from Bernoulli distribution with parameter p. Find suppose X 1, X 2, X 3, X 4, X 5 is a random sample from Bernoulli distribution with parameter p. Find suppose X 1, X 2, X 3, X 4, X 5 is a random sample from Bernoulli distribution with parameter p. Find suppose X 1, X 2, X 3, X 4, X 5 is a random sample from Bernoulli distribution with parameter p. Find suppose X 1, X 2, X 3, X 4, X 5 is a random sample from Bernoulli distribution with parameter p. Find suppose X 1, X 2, X 3, X 3, X 4, X 5 is a random sample from Bernoulli distribution with parameter p. Find suppose X 1, X 2, X 3, X 3, X 4, X 5 is a random sample from Bernoulli distribution with parameter p. Find suppose X 1, X 2, X 3, X 3, X 4, X 5 is a random sample from Bernoulli distribution with parameter p. Find suppose X 1, X 2, X 3, X 3, X 4, X 5 is a random sample from Bernoulli distribution with par Suppose X 1, X 2, X 3, X 4, X 5 for testing $H_0: p = 1/4$ against $H_1: p = 1/3$. What will be the conclusion if the BCR of size 0.1035 for testing $H_0: p = 1/4$ against $H_1: p = 1/3$. What will be the conclusion if the

BCK of characteristics are 1, 0, 0, 0, 1? BCK of servations are θ , θ as random sample of size 10 from Poisson distribution with mean θ . Obtain let S in θ is θ is θ in θ Let σ be size 0.0404 for the destriction of size 10 is drawn from N($\mu = 60$, σ^2) distribution. Obtain BCR of size 0.05 A random sample of $\sigma^2 = 2$ against $\sigma^2 = 4$. What will be the conclusion if the sample of $\sigma^2 = 2$ against $\sigma^2 = 4$. Arandom sample of size 10 is diamatical $H_1: \sigma^2 = 4$. What will be the conclusion if the sample observations for testing $H_0: \sigma^2 = 2$ against $H_1: \sigma^2 = 4$. What will be the conclusion if the sample observations for testing $H_0: \sigma^2 = 60.0, 61.2, 59.3, 59.7, 62.3, 56.5, 57.6, 53.4, 62.0?$ for testing 170. 61.2, 59.3, 59.7, 62.3, 56.5, 57.6, 53.4, 62.0? are 60.3, 60.0, 61.2, 59.3, 59.7, 62.3, 56.5, 57.6, 53.4, 62.0?

are 60.3, 60.0, 61.2, 59.5, 50.7, 52.5, 50.7, 53.4, 62.0? A sample of size n is taken from N(μ , $\sigma^2 = 25$) population. Obtain BCR for testing H₀: $\mu = 50$ A sample of size it is taken as $\mu = 50$. Also, find the value of n such that probabilities of type I and type II errors are against $H_1: \mu = 60$. Also, find the value of n such that probabilities of type I and type II errors are 0.025 and 0.01 respectively.

0.025 and 0.01 respectively. To test H₀: $\theta = 0.75$ against Arandom variable X has Poisson distribution with parameter θ . To test H₀: $\theta = 0.75$ against Arandom variable X has Poisson distribution with parameter θ . To test H₀: $\theta = 0.75$ against $\theta = 0.75$ a random sample of size 8 is taken. Obtain UMP test of $\theta = 0.75$ against A random variable 7. $\theta = 0.75$ against $\theta = 0.75$ against $\theta = 0.75$, a random sample of size 8 is taken. Obtain UMP test of size 0.1 assuming normal $\theta = 0.75$, a random sample of size 8 is taken. Obtain UMP test of size 0.1 assuming normal $\theta = 0.75$, a random sample of size 8 is taken. Obtain UMP test of size 0.1 assuming normal $\theta = 0.75$ against $\theta = 0.75$, a random sample of size 8 is taken. Obtain UMP test of size 0.1 assuming normal $\theta = 0.75$ against $\theta = 0.75$, a random sample of size 8 is taken. $H_1: \theta < 0.75$, and $H_2: \theta < 0.75$, and $H_3: \theta < 0.75$, and $H_4: \theta$ at 0 = 0.6, 0.5, 0.4, 0.3, 0.2, 0.1.

at $\theta = 0.0$, 0.0, p. a single observed value of x from a round in B to based on a single observed value of x from a round in a round in a round in a round Design Neymann 2 based on a single observed value of x from a population with probability distribution $H_1: p > 0.2$ based on a single observed value of x from a population with probability distribution $p(x) = q^{x-1} p$; x = 1, 2, 3,; (q = 1 - p, 0

; otherwise.

Assume that level of significance of the test is approximately 0.36. Evaluate the power function K(p) for p = 0.25, 0.30, 0.35, 0.40, 0.45, 0.50 and draw the power curve. A sample of size 10 is drawn from N(μ , σ^2 =10) distribution to test H₀: μ = 100 against A sample of Size 0.10. What will be the conclusion if the sample observations $H_1: \mu < 100$. Derive UMP test of size 0.10. What will be the conclusion if the sample observations are 100.1, 97.6, 133.5, 68.1, 82.3, 93.8, 102.5, 113.7, 79.2, 108.6? Also, find the value of power function when $\mu = 98$.

A random sample of size 9 is drawn from a population having p.d.f.

$$f(x, \theta) = \frac{1}{\theta} e^{-\frac{x}{\theta}}$$
; $x \ge 0, \theta > 0$
= 0 ; otherwise

Derive UMP test of size $\alpha = 0.05$ for testing $H_0: \theta = 2$ against $H_1: \theta > 2$. If a random sample from this population yields values 1.7, 3.5, 2.15, 3.1, 2.21, 1.78, 1.89, 2.17, 1.5; will the null hypothesis be accepted?

7. Y. B. Sc. Practical No. 7.2.3 LIKELIHOOD RATIO TEST (Derivations of the tests are not expected.)

Variable X – N (μ , σ^2). To test H₀: $\mu = 0$ agains: H₁: $\mu \neq 0$ where $\sigma^2 = 4$, what is the Let a region of size 0.05 specified by likelihood ratio test criterion? If the sample of size 10, you accept or reject H₀ at 5% level of significance? Let a random variable of size 0.05 specified by likelihood ratio test criterion? If the sample of size 10 gives 0.6, do you accept or reject H_0 at 5% level of significance? critical region of state of colors and the state of significance?

critical 166, do you accept follows Normal distribution with unknown mean μ and unknown standard the producer claims that on an average of 72 hours and a standard to the mean life of 72 hours and a standard to the mean life of 72 hours and a standard to the mean life of 72 hours and a standard to the mean life of 72 hours and a standard to the mean life of 72 hours and a standard to the mean life of 72 hours and a standard to the mean life of 72 hours and a standard to the mean life of 72 hours and a standard to the mean life of 72 hours and a standard to the mean life of 72 hours and a standard to the mean life of 72 hours and a standard to the mean life of 72 hours and a standard to the mean life of 72 hours and mean of a new battery and sample of 20 batteries showed the mean life of 72 hours and standard deviation of 12 hours. The producer claims that on an average a battery will last to a standard deviation of 12 hours and standard The life of of A random of of A random the producer claims that on an average a battery will last longer than 70 deviation of 12 hours. The producer claims that on an average a battery will last longer than 70 deviation of 12 hours and standard deviation of 12 hours. The producer of hours or producer's claim is true at 5% level of last longer than 70 deviation is the average life of size 10 is drawn from a Normal new true at 5% level of last longer than 70 deviation of the average life of battery from a Normal new true at 5% level of last longer than 70 deviation of the average life of battery from a Normal new true at 5% level of last longer than 70 deviation of the average life of battery from a Normal new true at 5% level of last longer than 70 deviation of the average life of battery from a Normal new true at 5% level of last longer than 70 deviation of the average life of battery from a Normal new true at 5% level of last longer than 70 deviation of the average life of battery from a Normal new true at 5% level of last longer than 70 deviation of the average life of battery from a Normal new true at 5% level of last longer than 70 deviation of the average life of battery from a Normal new true at 5% level of last longer than 70 deviation of the deviation of 12 hours of battery 70 hours or producer's claim is true at 5% level of significance?

hours, Is the average life of battery 70 hours or producer's claim is true at 5% level of significance? deviation is the average life of size 10 is drawn from a Normal population with mean $\mu = 70$ and unknown A random 2. In order to test H_0 : $\sigma^2 = 90$ against H_1 : $\sigma^2 \neq 90$, use likelihood ratio test hours is sample of size H_0 : $\sigma^2 = 90$ against H_1 : $\sigma^2 \neq 90$, use likelihood ratio test at 5% level of variance σ^2 and state your conclusion if the sample observations are variance σ . In other your conclusion if the sample observations are significance and state your conclusion if the sample observations are significant, 76, 64, 96, 72, 66, 77, 91.

A random sample of 2 and a standard deviation of 1.5 kg. Assuming weight loss follows Normal found to be 5 kg with a standard deviation of 1.5 kg. Assuming weight loss follows Normal found to be 3 kg the concluded that the population standard deviation of weight loss follows Normal distribution, can it be concluded that the population standard deviation of weight loss is 2 kg distribution, our distribution of distribution o

against the analysis and Y be independently distributed normal variables with parameters Let X and Y or X and X or X or X respectively where the common variance X is unknown.

 $N(\mu_1, \sigma^2)$ and $N(\mu_2, \sigma^2)$ and $N(\mu_1, \sigma^2)$ and $N(\mu_1, \sigma^2)$ and $N(\mu_1, \sigma^2)$ and $N(\mu_1, \sigma^2)$ and $N(\mu_2, \sigma^2)$ and $N(\mu_1, \sigma^2)$ and $N(\mu_2, \sigma^2)$ and $N(\mu_2, \sigma^2)$ are unknown. What is the likelihood ratio test to test $H_0: \mu_1 = \mu_2$ against $H_1: \mu_1 < \mu_2$ based on random samples drawn from the two populations? If the observed values are 26 27

31 28 Х

34 37 35 42 37 Y

What is your corclimion at 1% level of significance?"

A random sample of 7 brand X light bulbs yielded mean life of 891 hours and standard deviation of 10. A random sample of 10 brand Y light bulbs yielded mean life of 592 hours and standard deviation of 8. Use these data to test H_0 : $\sigma_1^2 = \sigma_2^2$ against H_1 : $\sigma_1^2 > \sigma_2^2$ at 1% level of significance. Assume that life of light bulb of both brands follow Normal law. Let 1.5, 2.4, 0.9, 1.3, 0.5, 2.6, 1.7, 0.4, 1.4, 0.8 be a random sample from Exponential distribution

with mean λ . Can you conclude that $\lambda \neq 2$ at 5% level of significance?

B.Sc. Practical No.7.2.4 NON-PARAMETRIC TESTS I

11 300

bulls is 350 hours. sample of female monkeys of a particular species are found to be as given below weights in a sample of s. 8.75, 8.40, 9.10, 9.25, 9.80, 10.05, 8.15, 10.00, 9.60, 9.80, 9.20 30, 9.50. Can it be concluded, using sign test at 1% level of significance, that the median 36, 8.43, 9.85 Can which the sample is drawn is greater than 8.41 kgs 2.41. 36, 8.43, 9.85. Can it be considered the sample is drawn is greater than 8.41 kgs.? (Use normal regulation.)

proximation.)

proximation.) centiles of 15 59 100 72 73 76 84 69 92 74 82 70 90 64 65 55 59 100 72 level of significance tes

64 65 55 level of significance to test whether the average percentile wilcoxon signed-rank test at 5% level of significance to test whether the average percentile services from this class is 72. (Use normal approximation.) of a student from this class is 72. (Use normal approximation.)

ore of a student from the salespersons in his insurance company spend at least 119 hours per manager claims that new salespersons in his insurance company spend at least 119 hours per manager claims that new salespersons revealed the following per field. manager ciairs and a random sample of 18 new salespersons revealed the following data.
onth in the field. A random sample of 18 new salespersons revealed the following data. onth in the new 1. 145 140 138 126 120 99 125 91 96 119 137 103 93 6 103 91 122 96 145 140 138 Wilcoxon signed and 1. 1 of significance. Using Wilcoxon signed and 1. 1 of significance.

103 91 122 137 103 93 113 5% level of significance, using Wilcoxon signed-rank test whether the data supports at 25% level of significance, using Wilcoxon signed-rank test whether the data supports of the normal approximation.) mager's claim. (Use normal approximation.)

mager's claum.

le following data shows the deviations from normal of daily temperatures in a certain city for two eks of a month.

 $\frac{20}{20}$ -0.6 5.3 -1 -2.3 0.7 -2.2 -0.9 2.9 3.4 2.5 5.2 5.8 -4.3nit be concluded that the pattern of deviations above or below normal temperature is a result of 1-random process? Use run test at 5% level of significance.

: following data of males(M) and females (F) was recorded during rush hours at a bus stop. run test at 5% level of significance to check if the arrangement of males and females in the ue is random. (Use normal approximation.)

NON-PARAMETRIC TESTS-II

Six randomly chosen models went on a diet in an attempt to lose weight (in lbs.) with the following results. following results.

Weight Before 115 136 133 128 153 135 139 131

Weight After Weight After Weight After Weight Lest at 5% level of significance to check if the diet is an effective means of losing

weight.
Test scores of a group of 15 high-school students before and after a training programme

are as given below.

are as given below. Score Before 63 78 58 70 75 76 74 88 74 94 50 95 94 84 86 75 97 72 100 101 98 98 Score After Score After Score After 1% level of significance to check if the training has any effect on the test Use sign test at 1% level of significance to check if the training has any effect on the test 105 84 90 scores, (Use normal approximation.)

Twelve children were asked to solve a puzzle under normal conditions and under the Twelve children and under the conditions of stress. The following data represents time(in seconds) taken to solve the

puzzle. 117 115 118 126 118 128 125 Normal 120 125 132 124 Condition 125 120 121 118 130 125 130 120 119 Stress 140 145 Condition

Do the data indicate sufficient evidence to conclude that time taken is more under conditions of stress? Use Wilcoxon test at 5% level of significance. (Use normal approximation.)

A Arandom sample of size 10 is obtained from one population and a random sample of size 11 is obtained from a second population. The observations are as follows.

Population- 0.26 1.00 1.29 5.00 8.00 16.00 1.60 8.20 10.20 12.81 I(X)

Pepulation- 0.16 0.08 0.10 0.00 2.02 0.80 2.56 1.28 1.26 5.10 6.24

Test, at 1% level of significance, using run test whether the population distributions represented by these two samples are different. (Use normal approximation.)

The average monthly expenditures (in Rs.) on cosmetics of working women (X) and housewives (Y) are listed below.

X 770 820 850 860 920 890 930 870 880 1000 910

Y 830 780 660 750 650 970 855 730 900

Use median test at 5% level of significance to check if there is a significant difference between the two sets of data.

The number of words in emails sent by boys (X) and girls (Y) were found to be as

X 85 89 86 91 .^Y 83 73 65 67 76 93 84 100 82 92 88

Use Mann-Whitney-Wilcoxon test at 5% level of significance to test whether the average number of words per email is same for boys and girls.

Twenty seven children, who enrolled for a swimming class, were divided into two groups. The children in group A were taught by traditional methods, while group B A and B. The children were taught by a new method. The instructor claimed that the new method is children were taught by a new method. At the end of the course, the awimming scores of the children in the two groups were as follows.

of the children in the two groups were as follows.

of the children in the second of t

TY.B.Sc. Practical No.7.2.6 SEQUENTIAL PROBABILITY RATIO TEST

Let X be a random variable having probability mass function (p.d.f.)1) Let X be a random variable having probability mass function (p.d.f.)

Derive SPRT of strength $\alpha = 0.05$ and $\beta = 0.10$ for testing H_0 : p = 0.5 against H_1 : pDerive SPR 1 of Section of the first sample observation has value 1? = 0.3 What will be the conclusion if the first sample observation has value 1? What will be N follows a binomial distribution (n = 10, p). Derive SPRT of A random variable N follows a binomial distribution (p = 10, p). Derive SPRT of $\alpha = 0.15$ and $\beta = 0.20$ for testing H_0 : $p = \frac{1}{4}$ against H_1 . A random variable $\beta = 0.20$ for testing H_0 : $p = \frac{1}{4}$ against H_1 : $p = \frac{3}{4}$. Strength $\alpha = 0.15$ and $\beta = 0.20$ for testing H_0 : $p = \frac{1}{4}$ against H_1 : $p = \frac{3}{4}$.

strength $\alpha = 0.15$ and accidents per day at a junction is assumed to follow a Poisson. The number of road accidents per day at a junction is assumed to follow a Poisson with unknown mean λ . In order to test $H_0 : \lambda = 1$ The number of Tourish with unknown mean λ . In order to test $H_0: \lambda = 1$ against $H_1: \lambda = 2$, it distribution with unknown mean λ . In order to test $H_0: \lambda = 1$ against $H_1: \lambda = 2$, it distribution with the distribution with the decision of the was decided to use SPRT of strength $\alpha = 0.10$, $\beta = 0.10$. What is the decision if the was decided to the decision if the was decidents observed on 8 days is 1, 1, 0, 2, 0, 1, 2, and 0 respectively? Also number of accidents observed on 8 days is 1, 1, 0, 2, 0, 1, 2, and 0 respectively? Also represent the test graphically.

represent the test μ = 0.5 distribution. Derive SPRT to test μ = 10 against 4) Let X follow N(μ , σ^2 = 0.5) distribution. Derive SPRT to test μ = 10 against 4) Let X ionow 10.7, $\beta = 0.05$, $\beta = 0.01$. What is the decision if the sample observations $H_1: \mu = 15$ with $\alpha = 0.05$, $\beta = 0.01$. What is the decision if the sample observations are 12.2, 12.9, 12.5, 13.2, 12.6, 11.9?

are 12.2, 12.3, 12.3, $(\mu = 120, \sigma^2)$ distribution. Derive SPRT to test $H_0: \sigma = 28$ against 5) Let X follow $N(\mu = 120, \sigma^2)$ distribution. Derive SPRT to test $H_0: \sigma = 28$ against Let A 1020. $\alpha = \beta = 0.05$. At what stage and what decision will be taken if the sample observations are 130, 185, 150, 165, 170, 200, 140?

6) A random variable X follows an exponential distribution with mean 1/0. Derive SPRT to test

 $H_0: \theta = 3$ against $H_1: \theta = 1$ with $\alpha = 0.20$, $\beta = 0.10$. What is the decision if the sample observations are 1.2, 0.5, 0.9, 2.3, 1.8?

aregies of the foreste where it is found to contain 195 hornest ones.

y.B.Sc. Practical No.8.1.1

GENETICS.J

The lack of pigmentation called albinism in man is the result of a recessive allele (a) The lack of pigmentation is a result of its dominant allele (A). Two normal parents and normal pigmentation child. Determine the probability that and normal prochild. Determine the probability that have an albino child is an albino. a) next child is an albino.

a) next two children are albinos.

b) the next to children one is albino and other is normal.
c) out of two children wire-haired to be insent gene W produces wire-haired to be insent gene wire-haired to be insent general to be insent gener c) out of two claims. We produces wire-haired texture in dogs; its recessive allele was smooth hair. A group of heterozygous wire haired in the state of the stat A dominant gone in A group of heterozygous wire haired individuals are crossed produces smooth hair. A group of heterozygous wire haired individuals are crossed produces smooth and their progeny are then test-crossed. Determine expected genotypic and and their progeny among test-crossed progeny phenotypic ratios among test-crossed progeny.

phenotypic ratios may be long (LL), round($\ell\ell$) and oval($L\ell$). If long are crossed shape of radishes may then F_1 are allowed to cross among them. Shape of Fausier of them F_1 are allowed to cross among themselves; what phenotypic with oval ones and then F_2 ?

ratios are expected in F2?

plack wool of sheep is due to recessive allele b and white wool due to its dominant Black Wood of the state of the allele D. A white lamb (male offspring), which is then backcrossed to a black. They produce a white lamb (male offspring), which is then backcrossed to a black. They produce the probability that backcrossing offspring being black? female parent. What is the probability that backcrossing offspring being black? temate partial partial delig bia An isolator. Yellow colour is governed by a dominant allele and white by its recessive allele. A random sample of 1000 kernels revealed that 910 were yellow. Find the allelic frequencies for this population.

The left handedness is assumed to be an effect of a single recessive trait. In a certain population, the prevalence of left handed people is 10%. If both parents are right handed, what is the probability that their first child is left handed? Baldness is governed by sex influenced trait which is dominant in men and recessive in women. In a sample of 10,000 men 6233 were found bald. In a sample

of equivalent size, how many non bald women are expected? The presence of horn in a special breed of sheep is dominant in males and recessive in females. If a sample of 900 female sheep is found to contain 225 horned ones

a) what percentage of females is expected to be heterozygous?

b) what percentage of males are expected to be horned?

A narrow reduced eye called 'bar' is dominant sex linked condition (B) in drosophila. Full wide type eye is produced by its recessive allele b. A homozygous wide type female is mated to a bar eyed male. Determine F1 and F2 phenotypic and genotypic expectations. (Assume F₁ are crossed among themselves to produce F₂). Among New-York Caucasians, the frequency of ABO blood groups were found 49% type O, 36% type A, 12% type B and 3% type AB. Obtain allelic frequencies using

a) Bernstein's method

b) gene counting method.

GENETICS-II

F.B.Sc. Practical No.8.1.2 masummer-squash, white fruit colour is governed by a dominant allele W and may by its recessive allele w. A dominant allele at another independent In a summer-squasity with the disc shaped variety in produces disc shaped variety in the disc produces disc shaped fruit and its recessive allele d yields sphere shaped by the Homozygous white disc shaped variety is crossed with homozygous who produce F. pellon produces and the disc shaped variety is crossed with homozygous yellow full meshaped variety to produce F1. full named variety to produce F₁. sphere shaped variety are allowed to cross at random among themselves, obtain phenotypic ratios a) If F1 are allowed to cross at random among themselves, obtain phenotypic ratios

in Figeneration.

In Figeneration what phenotypic ratios are expected in F₂?

In Figeneration in F₂? b) If F_1 are test closses governed by a dominant allele L and long by its recessive short hair in rabbits is governed by a dominant genotype D short hair in rapidle 1. Black hair results from the action of dominant genotype B and brown allele 1. Black hair results from the action of dominant genotype B and brown allele 1. Black hair results from the action of dominant genotype B and brown allele l. Black in genotype bb. Determine expected genotypic and phenotypic ratios in from recessive genotype bb. the progeny from the cross

the program Lebb b) LeBb × LeBb c) dihybrid testeross.

a) Libbox De a yellow cotyledon is dominant to green and inflated pod shape is in a garden pea, yellow cotyledon. When both traits are considered in In a garden pour stricted form. When both traits are considered jointly in self dominant to constricted form anneared in the following dominant to domina fernized and 193 green and inflated, 556 yellow and inflated, 184 yellow and constricted, 184 yellow and inflated, 184 yellow and constricted. Test the data for independent assortment.

The shape of radish may be long (LL), round ($\ell \ell$) or oval ($L\ell$) and its colour may le red (RR), white (rr) or pink (Rr). If long white strain is crossed with round red strain, what phenotypic ratios are expected in F₁ and F₂? (F₁ are crossed among

themselves to produce F2)

Agene locus with co-dominant alleles is known to govern feather colour in chickens such that BB- black, Bb- blue, bb-splashed white. An independently segregating locus L governs leg length such that LL-normal length, L \ell-squatty type leg 11- genotype is lethal. Determine phenotypic ratio among the matured progeny if dihybrids are crossed among themselves.

A dominant allele Y is responsible for yellow seed colour and its recessive allele y produces green. Co-dominant allele at independent assorting locus R specify the shape, such that RR-round, Rr-oval and rr- shrunken. From a cross between yellowoval and green-oval, obtain the phenotypic ratios expected in the progeny.

Adominant allele L governs short hair in guinea pigs and its recessive allele l governs long hair. Co-dominant allele at an independently assorting locus specify hair colour. YY-yellow, Yy- cream, yy-white. From crosses between dihybrid short-cream pigs, the following phenotypic distribution is observed. 50-short& cream, 21-short & yeilow, 23-short & white, 21-long & cream, 7-long & cream and 6-long & white. Are data consistent with Mendel's laws?

by the was assumed C stands for colour and R (independent of C) is responsible for red colour, with recessive allele r responsible for yellow. cc represents white (c is recessive). From dihybrid crosses in the offspring generation; out of 160 plants; 90 showed red colour, 25 yellow and rest white. Test the data for recessive epistasis.

Coloured alueron in corn is due to interaction of two dominant genes in the Coloured alueron in the genotypes at this locus produce colouriess alemone. A genotype A C; all other genotypes at this locus produce colouriess alemone. A genotype A C; all other genotypes at this locus produce colouriess alemone. A genotype A C; all other genotypes at this locus produce colouriess alemone. A genotype A C; all other genotypes at this locus produce colouriess alemone. genotype A C, an one is test-crossed to pure colourless strain, the F₁ are with homozygous coloured. When F₁ are crossed among thems lives to produce F₁ homozygous coloured. When F₁ are crossed among thems elves to produce F₂, W₂ coloured alcurone. Went 2460 colourless. Test the data for dusting 3300 coloured & 2460 colourless. coloured aleurone. 2460 colourless. Test the data for duplicate recessive exhibits 3300 coloured & 2460 colourless.

In an experiment on all 25, 275, 975 and 625 respectively. Obtain allelic asB_asies.

frequencies.

GENETICS-III

In a dihybrid testeross it was observed that the phenotypes A_B_, A_bb, aaB_ and aabb occurred 67, 19, 29 and 77 times respectively. Verify whether there is linkage, if so obtain measure of linkage by m.l.e. method, also obtain its S.E. (Assume genes are in coupling phase)

In dihybrid intercross (genes are in coupling phase) the observed frequencies of the phenotypes R_Y_, R_yy, rrY and rryy in F₁ generation were 187, 35, 37, 31 respectively. Test whether there is a linkage, if so obtain measure of linkage by m.l.e. method, also obtain its S.E.

In dihybrid cross AB/ab × AB/ab the observed frequencies of the phenotypes A B, A bb, aaB and aabb in F₁ generation were 1221, 219, 246, 243 respectively. Estimate the recombination factor between A and B by

a) square root method

b) product ratio method

In a cross of the type Ry/rY × Ry/rY the following results were found.

Phenotype R.Y. R.yy rrY rry Observed freq 360 120 160 20

Estimate the recombination factor between A and B

a) square root method

b) product ratio method

EPIDEMICS-1

- For a simple epidemic with one infected person in a population of size 86 and the infection rate as 0.35, draw the epidemic curve by taking τ = 0.03, 0.04, 0.05, 0.06, 0.07, 0.08. When the curve attains maximum? What are the number of susceptibles and number of infectives at that time?
- In a simple epidemic there were 55 susceptibles and 3 infectives at the start. The infection rate is 0.95, $\tau = t\beta$ obtain values of ω when $\tau = 0.01$, 0.02, 0.03, 0.04. 0.05, 0.06, 0.07, 0.08, 0.09. Draw the epidemic curve. At what value of τ epidemic curve attains its maximum? What are the number of susceptibles and infectives at that time?
- In a general epidemic model with removal, x and y respectively represent the number of susceptibles and the number of infectives at any time t. If the total size of the group is 500 with x0 = 450 and relative removal rate is 300, obtain values of y when x take the values 450, 400, 350, 250, 200,150. What is the maximum value of y? At that time how many individuals are expected to be removed? According to threshold theorem what will be lower limit of x?
- 4) Assuming that an epidemic has started with five carriers in a group of 200 susceptibles, find an estimate of the total size of the epidemic if the relative removal rate for carriers is 4.
- 5) For a carrier model there are 275 susceptibles and 17 carriers at the start, infection rate is $\beta = 0.1$ and removal rate is $\gamma = 0.375$. Find the number of susceptibles that will escape infection.
- 6) For a carrier model with 20 carriers and 450 susceptibles at start, infection rate is $\beta = 0.0075$ and removal rate $\gamma = 0.075$, find the number of susceptibles left uninfected at the end of 5 days, 10 days, 15 days and 20 days. Will the situation change substantially if by intensive drive the removal rate for carriers is doubled?

T.Y.B.Sc. Practical No.8.1.5

EPIDEMICS-II

- The following data is collected from households of size 3 with one primary case of The long train one primary case of meesles. Estimate the value of p, the chance of adequate contact and obtain its S.E. $\{1,1\}$ $\{1,1,1\}$ Type of chain {1} 25 135 Observed freq 34
- Estimate the value of p, the chance of adequate contact between a susceptible and an Estimate in conjunctivitis when 500 households each of size 3 with single infective in conjunctivitis when 500 households each of size 3 with single introduction were considered. It was reported that in 180 households the disease spread to two members and in 197 household all members are affected. Also obtain its Standard error.

Assuming all households are of size 4, estimate p, the chance of adequate contact. The observed frequencies of the chains {2}, {2,1}, {2,1,1} and {2,2} are 123, 180, 73 and 124 respectively. Use a) Greenwood model

b) Reed Frost model

- Estimate p-the chance of adequate contact between susceptible and infective in an epidemic from a data of 600 households each of size 4, started with two introduction, only two members were infected in 143 households whereas all members were infected in 217 houses. Use Greenwood model.
- 5) Estimate p-the probability of adequate contact between a susceptible and infective. Data is collected from 175 households each of size 4 with single introduction. The observed chain frequencies of the chains {1}, {1,1}, {1,1,1}, {1,2}, {1,1,1,1}, {1,1,2}, {1,2,1} and {1,3} are 10, 9, 6, 20, 11, 8, 20 and 116 respectively. Estimate p using
 - a) Greenwood model

b) Reed Frost model.

- 6) For a household of size 4, find the probabilities of the chains {1,2,1} and {2,1}, $\{2,1,1\}$ using
 - a) Greenwood model

b) Reed Frost model

assuming probability of adequate contact between susceptibles and infective is 0.05.

T.Y.B.Sc. Practical No.8, 1,6

BIOASSAYS-I

- 1) Lethal doses of two preparations A and B required for a fixed response are given to Lethal doses of the seven experimental animals each. Obtain the relative poteny two difference to B assuming normality of of A with reference to B assuming normality of
 - a) doses 8
 - b) log withms of doses
 - 1.89 2.34
 - A 1.55 1.58 1.71 1.44 1.24 B 2.42 1.85 2.00 2.27 1.70 1.47 2.20
 - Also obtain 95% fiducial limits for the relative potency in both the cases.
- Also obtains data shows the amount of drug A and drug B required to cause cardiac
 - arrest in pigs. 46.0 46.7 44.9 41.0 45.2 50.4 36.9 40.0 Dose A mg/kg
 - Dose B mg/kg 68.6 63.3 73.6 68.0 69.1 65.5 64.4 65.0
 - Eight pigs were used for each drug. Using log dose as metameter, estimate the potency of B with respect to A as standard preparation. Also find the 95% fiducial limits for the same.

y.B.Sc. Practical No.8, 1.7

BIOASSAYS-II

In a (2.2) point parallel line assay, dosages were $T_1 = 28.0$ mg, $T_2 = 42.0$ mg of test (2.2) point $T_2 = 42.0$ mg of test preparation $T_1 = 28.0$ mg, $T_2 = 42.0$ mg of test preparation $S_1 = 14.0$ mg, $S_2 = 21.0$ mg of standard preparation

Responses were given as follows. T_2 T_1 S_2 65 46 Sı 62 72 53 43 62 74 58 48 66 76

52
Estimate the potency of test preparation with standard preparation after verifying
Estimate the potency obtain 95% fiducial limits for the notency Estimate the Policy obtain 95% fiducial limits for the potency.

owing responses were observed in a (3,3) point assay of vitamin B₁₂.

llowing respons	Sı	S ₂	S ₃	T_1	T ₂	T_3
Dose	0.8	1.2	1.8	4	6	9
D. monge	65	75	86	60	78	84
Response	60	76	83	62	73	84
	61	68	83	67	66	83
	45	55	82	65	75	85
7, 11	72	75	82	58	73	79
and the f	62	71	84	70	71	84

Estimate the potency of the test preparation relative to standard preparation after canying out the necessary validity tests.

In a (3,3) point-parallel line assay, doses were $S_1 = 0.12$, $S_2 = 0.24$, $S_3 = 0.48$, and $T_1 = 0.25$, $T_2 = 0.5$, $T_3 = 1.00$. (with usual notations)

Responses were given as follows.

Dose Dose	S1	S2	S3	T1	T2	T3
Response	298	250	230	274	176	19
Acceptance	280	193	228	293	231	17
. **	262	228	199	325	222	20
	304	259	216	339	189	18
	300	256	201	301	210	20

Test for assay validity.

y.B.Sc. Practical No.8, 1,8

BIOASSAYS-III

Eight doses of a preparation of insulin were injected to batches of mice and the Eight doses of a proposition of collapse was recorded. Table below gives number of mice showing the symptoms of collapse was recorded. Table below gives number of mice should be stated (n) and number of convulsed mice(r). Using dose (z), the number as a dose metameter estimate LD 50 for at dose (z), the number of the dose as a dose metameter estimate LD 50 for this preparation using logarithm of the dose as a dose metameter estimate LD 50 for this preparation using

logarithm of the		en s					
anabit filedit	7.0	8.5	10.5	13	18	21	28
$\frac{z}{32}$	38	37	40	37	31	37	30
$\frac{n}{5}$	11	14	18	21	23	30	27
1 1			· 1 /-1-		-114		4

te the relative potency using probits. Make natural log transformation.

ate the relati	OARD PREE	ARATION
Z	n	seed the same
• 3.4	33	2
5.2	32	5
7.0	38	11
8.5	37	14
10.5	40	18
13. 0	37	21
18.0	31	23
21.0	37	30
28.0	30	27
1 7.0 . 0	100	

PREPARA	ПОП
n	r
40	2
30	10
40	18
35.	21
47	27
	n 40 30 40 35

The following table gives the numbers of frogs tested (n) with different doses (z) of a drug and the number of survivals (r). Log transformation ($x = log_{10} Z + 4$) is used for linearity. Estimate LD50 using probit method.

20	20	20	18	10	10
7 to 18 h		11 11 2			
20	16	07	04	01	00
100	25	250	300	350	400
	75 0 140 - 1 150	20 16	20 16 07	20 16 07 04	20 16 07 04 01

T.Y.B.Sc. Practical No.8.2.1

INVENTORY MODELS-I.

A shopkeeper has a uniform demand of an item at the rate of 600 items per year. He

A shopkeeper has a uniform demand of an item at the rate of 600 items per year. He

Buys from a supplier at a cost of Rs.8 per item and the cost of ordering is D. 10 A shopkeeper has a cost of Rs.8 per item and the cost of ordering is Rs.12 each bys from a supplier at a cost of Rs.8 per year of value, how from the stock holding costs are 20% per year of value, how frequency should he time the stock and what is the optimal order quantity? What is the ime. If the stocks and what is the optimal order quantity? What is the minimum eplenish his stocks and what is the optimal order quantity? What is the minimum eplenish cost?

total relevant control of a machine for its annual A manufacturing company purchases 18,000 parts of a machine for its annual A manufacturing one month's usage at a time. Facilities of the same of the control of the A minimacrimus one month's usage at a time. Each part costs Rs.20. The requirements, ordering one rooter is Rs.30 and the carrying charges are 150. requirements, or costs Rs. 20 and the carrying charges are 15% of the average ordering cost per year. Is the current ordering policy the best for the second costs of the average of the second costs are 15% of the average of the second costs are 15% of the average of the second costs are 15% of the average of the second costs are 15% of the average of the second costs are 15% of the average of the second costs are 15% of the average of the second costs are 15% of the average of the second costs are 15% of the average of the second costs are 15% of the average of the second costs are 15% of the average of the second costs are 15% of the average of the second costs are 15% of the average of the second costs are 15% of the average of the second costs are 15% of the average of the average of the second costs are 15% of the average of the second costs are 15% of the average of the second costs are 15% of the average of the 15% of the average of the 15% of the ordering cost per of the average of the period policy inventory per year. Is the current ordering policy the best for the company? If not, inventory per year. Is the current ordering policy the best for the company? inventory per your policy? What would be the company's annual saving if it what is the optimal policy?

adopted the optimal policy?

The ABC company sells its product at the rate of 500 units per day throughout a The ABO waits per day. To satisfy this demand the company manufactures it at the working year and solve of machinest and other and other material, Re.1.00 for labour and Rs.2.50 for hire of machinery and other variable overheads. The setup cost per batch would be Rs.600 and the stock-holding costs have been estimated at 20% per annum. Calculate the optimum quantity to be manufactured in each batch and the total inventory cost for the year.

Suppose you are reviewing the production lot-size decision associated with a production operation where the production rate is 8000 units per year, annual demand is 2000 units, set-up cost is Rs. 300 per production run and holding cost is Rs.1.60 per unit per year. The production run is 500 units every 3 months. Would you recommend a change in production lot size? If so why? How much could be saved, if any, by adopting the new production lot size?

A dealer supplies you the following information with regard to an item of inventory.

Annual demand:

5000 units.

Ordering costs:

Rs.250 per order

Inventory-holding costs.

30% of the value of inventory per year

Inventory stock-out costs:

Rs.10 per unit per year

Price:

Rs.100 per unit.

Find

a) Economic order quantity.

b) What quantity should be allowed to be backordered?

() What will be the maximun: inventory at any particular time of the year?

d) Optimum total relevant cost.

A furniture dealer sells special computer chairs. Each purchase order costs Rs.50 to the dealer and the holding costs amount to Rs. 80 per chair per year. The dealer sells

% chairs per month. He has estimated the shortage cost at Rs.20 per chair per year. What is the EOQ for the chair?

b) When the supply of the chairs comes, how many chairs are expected to be

customers immediately?

What is the minimum total relevant cost?

A company has a demand of 12000 units/year for an item and it can produce 2000 such items per month. The cost of one setup is Rs.400 and the holding cost per unit per month. The shortage cost of one unit is Rs.20 per year. Find the optimum lot size and the total cost per year assuming Rs.4 as the cost per unit. Also find the maximum inventory level.

T.Y.B.Sc. Practical No.8.2.2 INVENTORY MODELS-II A magazine vendor buys magazines from its publisher for Rs.10 each and sells them A magazine venue. The magazine is a weekly magazine and the probability distribution for Rs. 15 each. The magazine as follows. for Ks. 1. demand is given as follows.
of weekly demand 30 31 33 34 36 Demand: 0.09 0.13 0.18 0.18 0.15 0.12 0.15 probability: probability: 0.15
probability: 0.15
probability: 0.15
if the vendor cannot return unsold magazines how many magazines should the vendor order each week? vendor order cache vendor order each product is Rs.1,000 and its holding cost per the purchase price of a unit of a certain product is Rs.1,000 and its holding cost per period is Rs.100. If the order quantity is 4 units, find the provider period is Rs.100 and its holding cost per The purchase price of the order quantity is 4 units, find the permissible range unit per period is Rs. 100. If the order quantity is 4 units, find the permissible range of C2 given the following probability distribution of demand. Demand: Demand.

Probability: 0..05 0.10 0.10 0.20 0.25 0.15 0.05 0.05 0.05

Probability: opposition of the probability of the probabil A sweet-meat company sells its 'Mango Barfi' by weight. The barfi fetches the best A sweet-mean day it is prepared but there is an unlimited market for one day old barfi. The company makes a profit of Rs.20 per kg. on fresh barfi but a loss of Rs.4 for one The company old barfi. The daily demand for the product is a random variable with p.d.f. given below. f(x) = 0.1 - 0.005 x; $0 \le x \le 20$; otherwise How many kgs. of the barfi must the company make daily? A wholesaler in flowers wants to decide how many baskets of flowers he must procure every morning. He estimates the demand for flowers to be rectangular between 30 and 40 baskets. The purchase price for one basket is Rs 50 and the selling price is Rs.80. One day old flowers are worthless. What would be your advice regarding the number of baskets the wholesaler must procure every morning? The probability distribution of weekly sales of a certain item is as follows. 2 1 3 4 Weekly sales: 6 0.03 0.06 0.15 0.34 0.23 0.12 0.07 Probability: The cost of carrying one unit for one week is Rs.40 and the cost of understock is Rs.60 per unit per week. If the demand occurs at a uniform rate over the week how many units of this item must be stocked at the beginning of each week? The demand for 'integrated circuits' at an electronic goods dealer is a random variable with probability distribution as given below. Weekly demand(in '00 units): 0 1 3 4 5 6 Probability: 0.08 0.10 0.22 0.25 0.15 0.15 0.05

Determine the optimum inventory policy for the dealer if he is to order the circuits from a manufacturer at the beginning of each week. From past experience he has found that the cost of keeping 100 circuits for a week is Rs.0.20 while the loss due to not having the 100 circuits in stock is Re.1 per week. Assume demand occurs at a uniform rate.

the daily demand in kgs. for a certain product is a continuous random variable with

f(x) = 0.05; $0 \le x \le 20$

the cost of making one kg. of this product is Rs. 180 while the selling price is Rs. 300 kg. Hour and occurs at a uniform rate? How many kgs. must be made daily if the demand occurs at a uniform rate?

The daily demand for a certain commodity is a random variable with a probability The daily demand a rectangular between 500 and 1,000 units. The cost of distribution which is rectangular between 500 and 1,000 units. The cost of distribution which a unit is Rs. 80. If the inventory policy is to stock 700 units every manufacture of a unit is Rs. 80. Assume that all uncold units manufacture of a data selling price. Assume that all unsold units at end of day are

worthless.

The ABC company is engaged in the manufacture of personal computers. The XYZ

The ABC company is engaged in the manufacture of personal computers. The XYZ The ADC company undertakes to supply 24000 chascis every year to the computer company. Company unto the company being a small-scale entrepreneur undertakes to supply the The XYZ company being a small-scale entrepreneur undertakes to supply the The A12 country of a daily basis. It finds that the cost of holding a computer chassis chassis required on a daily basis. It finds that the cost of holding a computer chassis chassis required and there is a clause in the contract penalizing in a stock is Re.1 per unit per month and there is a clause in the contract penalizing in a Stock is the chassis at the rate of 2000. it Rs. 2 per that the chassis at the rate of 3000 chassis per month and the setup company produces that the Determine cost is Rs.3500 per batch. Determine

a) How frequency should batches be started and what should be the batch size?

The state of

b) What is the minimum total relevant cost? b) Wild is the Annual demand for a particular item of inventory is 10000 units. Inventory carrying Annual definition of the state by the supplier is Rs.4 per unit. However the supplier is willing to give a discount of 5% for orders of 1500 or more. Is it worthwhile to avail of discount offer? of 570 for optimal order quantity for a product for which price breaks are as follows:

Quantity ordered (units)	Unit cost (Rs.)			
$0 \le Q < 100$	20			
$0 \le Q < 100$ $100 \le Q < 200$	18			
	16			
200 ≤ Q	. TT			

The monthly demand for the product is 400 units. The storage cost is 20% of the

of the product and the cost of ordering is Rs.25 per order.

The annual demand for a product is 64,000 units. The ordering cost is Rs.10 per order and the estimated cost of carrying one unit in stock for a year is 20%. The normal price of the product is Rs.10 per unit. However, the supplier gives a quantity discount of 2% on an order of at least 1,000 units at a time and a discount of 5% if the order is for at least 5,000 units. Suggest the most economic purchase quantity

Company A wants to know what production cost its major competitor Company B per order. has assigned to product item X. After a bit of investigation Company A has collected the following data about Company B's production process:

Production lot size......2,600 units Annual demand......30,000 units

Company A has further learnt that Company B produces according to the economic lot size model. What is the Company B's cost of producing product item X? (1 year = 300 days).

T.Y.B.Sc. Practical No.8.2.3 REPLACEMENT THEORY

is considering replacement of a machine, whose cost is Rs. 12, 200 and the scrap value (maintenance & operating) costs are found to be as follows.

cont is aming	Mann	011111		5	7 00013	are rou	na to b	e as lon
Rs. 200. The running	1	2	3	4	5	6	7	8
Rs. 200	200	500	800	1200	1800	2500	3200	4000
Year: Running cost (Rs.): Running should the mac. When should the mac.	nine be	repla	iced?				rately.	21/2 11/2
when should the flac	unning	g a ma	cline,	the cos	st of wh	nich is R	cs.60,00	00, are g

1)

the tracted in running a machine, the cost of which is Rs.60,000, are given below

_	1 0			
1	2	3	4	5
42,000	30,000	20,400	14,400	9,650
4,000	4,270	4,880	5,700	6,800
14,000	16,000	18,000	21,000	25,000
	4,000	4,000 4,270	4,000 4,270 4,880	4,000 4,270 4,880 5,700

Determine, the optimum period for replacement of the machine.

Determine, and or Company owns three mini buses, two of which are two years old while the The Mini Transport Company owns three mini buses, two of which are two years old while the The Muni Transport of these buses was purchased for Rs.8 lakhs. The company did one is only a year old. Each of these buses by two full-sized buses. dird one is only like three buses by two full-sized buses, each such bus containing 50% contemplates replacing the three buses by two full-sized buses, each such bus containing 50% contemplates are stated by the Full- sized by the Full- sized by the following data, state whether the mini buses be replaced by the Full- sized buses. If not, state why? If yes, state when?

	and the state of	1.01	CILIVIII	DOS				
Year	1	2	3	4	5	6	7	8
Running cost(Rs.000's)	30	36	48	50	80	112	150	200
Resale value (Rs.000's)	700	610	550	490	320	200	100	50
Kesare varia	nere est	FOR A I	TULL-SIZ	ZED BUS	3	Caralter of a		
Year	1	2	3	4	5	6	7	8
Running cost(Rs.000's)	34	39	47	58	72	90	120	160
Resale value (Rs.000's)	1000	920	860	810	760	660	540	400

The data on the running costs per year and resale price of equipment A whose purchase price is Rs.200, 000 are as follows.

Year	1	2	3	4	5	6	7
Running cost(Rs.)	30,000	38,000	46,000	58,000	72,000	90,000	110,000
Resale value (Rs.).	100,000	50,000	25,000	12,000	8000	8000	8000

2) What is the optimum period of replacement?

When equipment A is two years old, equipment B which is a new model for the same usage is available. The optimum period of replacement is 4 years with an average cost of ks.72,000. Should equipment A be changed with equipment B? If so, when?

Wo wheeler is costing Rs. 50, 000 and operating & maintenance cost for its use over a specific Is costing Rs. 50, 000 and operating & manufacture are estimated to be Rs. 8000 for each of first five years and increase by Rs. 2000 per year for sixth year onwards. If money is worth 10% per year, when should the two wheeler be splaced optimally? (No scrape value)

Amanufacturer is offered two machines A & B. The price of A is Rs. 5000 and running costs are Marked at Rs. 800 for the first five years each, increasing by Rs. 200 every year after fifth year. White B, which has similar capacity as A, costs Rs. 2500, but will have running costs of All of the first six years, increasing by Rs. 200 per year thereafter. If money is worth Poply year and there is no scrap value for the machines, determine which machine should be

ARUN company limited provides ferry service across the Ganges. One of their boats is in poor condition and requires immediate repairs ferry Rs. 20.000. The ferry however can also be disposed off right new for 7) costing Rs. 2000. If repaired and used it will have a life of ten years but will need minor Rs. 7000. If repaired of five years costing Rs. 8000. Operating cost of the ferry will be repairs at the cost of the ferry will be repairs are carried out. At the end of ten years, the Rs.16000 each years, 16000 each years, 16000 each years, ferry will have scrap value of Rs. 5000. As an alternative, the company may ferry will have a new ferry for Rs. 40, 000. It has an estimated life of ten years with a purchase a new purcha minor repair will be Rs. 12000 every year. At the end of ten years the ferry will cost for the ferry value of Rs. 6000. What should the company do is cost for the value of Rs. 6000. What should the company do if money is worth 12% have a solution operating cost incurred at the beginning of the year) per year. (Asystem contains 10,000 resistors each of which is to be replaced on failure. The A system of replacing an individual resistor is Rs. 1, while if all resistors are replaced at same time, the cost per resistor would be reduced to only 35 paisa. The percentages of resistors surviving S(t) at the end of month 1 are as follows.

Design the optimum replacement policy.

A large computer installation contains 2000 components of identical nature which are subject to failure as follows.

 Month end
 1
 2
 3
 4
 5

 % failure to date
 10
 25
 50
 80
 100

The cost of individual replacement is Rs.3. If all the components are replaced in one lot, then the cost per component is Rs.1.

- a) Assess which policy of replacement would be economical.
- b) If group replacement is economical at current costs, then assess at what cost of individual

replacement would be uneconomical.

c) How high can the cost per unit in group replacement be to make a preference for individual

replacement policy?

10) In a factory 1000 lamps are in use. The life distribution of lamps is as follows.

 Life (in hours)
 300 - 400
 400 - 500
 500 - 600
 600 - 700
 700 - 800

 Probability
 0.07
 0.13
 0.35
 0.27
 0.18

Cost of replacement on failure is Rs.6 per lamp. Cost group replacement is Rs.1.8 per lamp. Assuming that failures occur just before the end of an interval, whenever they occur during an interval. Determine optimum replacement policy.

DECISION THEORY

No.8.2.4 Adealer in second hand machinery is offered 5 machines by a company for Rs. 5000/- only. He Advaler in second hand the machines for Rs.2200/- at a fair, but he knows that any machines not sold expects to sell each of the machines for Rs.2200/- at a fair, but he knows that any machines not sold expects to sell each of the machines as many machines as he call A dealer to sell each of the dealer has the option of buying as many machines as he chooses, at the rate of D. where the various principles of choice, determine the control of D. where the assumption of the dealer has the various principles of choice, determine the control of D. where the cont expects to waste and not so so waste and not so so waste and not so would be a waste and not so would be a waste and not so would be a waste and not so so would be a waste and not so so would be a waste and not so so waste and not so waste an would be dealer has the options principles of choice, determine the optimal strategy for the dealer in hease. Using matrix, the elements of which indicate the

h case.

In the following matrix, the elements of which indicate the costs, obtain the decisions using an the following of decision making. oinles of decision making.

From minciples	01	-	_
airent plus	TA	Δ.	1
from a principles	A ₂	713	-

TA.	A ₂	A ₃	14	A15
e. 26	22	13	22	18
S ₁ 26	22	34	30	20
S ₂ 20	22	18	18	20
$\frac{ S_3 }{ S_3 } \frac{10}{22}$	22	18	18	18
15,12-1				

Anewspaper boy is thinking of selling a special one time edition of a sports magazine to his regular an customers. Based on his knowledge of his customers, he believes the A newspaper customers. Based on his knowledge of his customers, he believes that he can sell henveen 9 to 12 copies. The magazines can be purchased at Rs. 8/- each and can be sold for Rs. 12/-Magazines that are not sold can be returned to the publisher for a refund of 50%. Construct the payoff matrix for the above problem.

Constitute of the control of the probabilities associated with selling of 9 to 12 copies are 0.30, 0.40, 0.20, 0.10 respectively, determine the best decision using the E.M.V. approach. of Find the E.P.P.I. and hence find the E.V.P.I.

Amotor parts dealer finds that the cost of a particular item in stock for a week is Rs. 10/- and the and of a unit shortage is Rs.30/-. The probability distribution of weekly sales (in '000) is as follows.

Weekly Sales (in '000):	0	1	2	3	4	5	6
Probability:	0.10	0.10	0.20	0.25	0.15	0 10	0.10

How many units per week should the dealer order? Also find the E.P.P.I.(Expected payoff (cost) with perfect information) and hence the E.V.P.I.

For the past 200 days, the sales of bread from the Fresh Food Bakery have been as follows.

Daily Sales (loaves):	0	100	200	300	400
No. of Days:	10	60	60	50	20

The bakery's production costs are Rs.2 per loaf, sale price is Rs.4 per loaf, and any bread unsold at the end of the day is contracted to a local farmer who pays Re.1.per loaf. Draw up a Myoff table for each sales/production combination. Obtain the regret matrix.

Determine the best action using the E.O.L. approach.

Malis the maximum amount the bakery owner should be willing to spend for perfect Mormation abut the number of loaves expected to be demanded in a day.

A farm owner is considering drilling a well. In the past only 70% of the wells had yielded water at a A farm owner is continued area. On finding no water at a depth of 200 feet some persons drilled depth of 200 feet in the area. On finding no water at a depth of 200 feet some persons drilled depth of down upto 250 feet but only 20% struck water at that depth. The cost of depth of the door water down upto 250 feet but if he door water that if he door water that if he door water at that depth. depth of 200 feet in the deep the down up to 250 feet but only 20% struck water at that depth. The cost of drilling is Rs.50 per further down up to 250 feet but only 20% struck water at that depth. The cost of drilling is Rs.50 per further farmer estimates that if he does not have his own well he will have to pay Be 15 Rs.50 per further down upto 25 that if he does not have his own well he will have to pay Rs. 15,000 over foot. The farmer estimates that if he does not have his own well he will have to pay Rs. 15,000 over foot. 10 years. (in Pv terms) to buy water from a neighbour. foot. The faither (in Pv terms) to buy water from a neighbour, the next 10 years. (in Pv terms) to buy water from a neighbour. The following decision could be optimal.

a) Do not drill a well.

b) Drill upto 200 feet. b) Drill upto 250 feet, drill further upto 250 feet.

c) It no control of the control of t E.M.V. approach.

E.M. T. A manufacturer of a popular brand of refrigerator intends to market its new fuel economy model. It has three options as follows.

a) To go ahead and launch.

b) To run a test market.

c) To abandon the model.

If the product is launched, the demand may be poor, good or excellent. The probabilities of poor, If the product is the product is used at a cost of good or excellent demand are 20%, 40% and 40% respectively. If a test market is used at a cost of good or exceeding give the indication of market being either favourable or unfavourable, there being Rs.2 lakhs, it will give the indication of market being either favourable or unfavourable, there being RS.Z IANID, 12 Unitary of unitary ourable, the 50:50 chance of market being favourable and unfavourable. If the test market response is favourable and the product is launched, the probability of poor, good and excellent market are 10%, 1avourable and the product is 20% and 70% respectively. And if the test market response is unfavourable and the product is Junched, the corresponding probabilities of poor, good and excellent market are 80%, 10% and 10%. The payoffs of the above categories of demand are Rs. 4 lakhs, Rs. 8 lakhs and Rs. 14 lakhs

a) Ignoring the expense already incurred, draw a decision tree for the above problem.

b) Suggest an optimal strategy for the manufacturer.

Alandowner has an option to sell rights for oil exploration. If he sells the rights, he can get Rs.5 lacs and a further Rs.60 lacs if oil is found. However, as an alternative he can himself conduct a test to check whether oil is there, the cost of which is Rs.12 lacs. If the test succeeds, he can expect a profit of Rs.120.lacs by selling the oil. Alternatively, the landowner can have soundings performed at a cost of Rs.3 lacs. The test of soundings is perfect 70% of the times if oil exists and 90% of the times, if oil does not exist. If it is expected that the oil exists with a probability of 0.3, what should the landowner do? Use a decision tree.

THEORY OF GAMES

Identify the saddle point in the following payoff matrices and the pure strategies for each player:

$$A \begin{bmatrix} 7 & 3 & 4 & 10 \\ 3 & 0 & -2 & -4 \\ 6 & 6 & 6 & 7 \\ -3 & 4 & 2 & 3 \end{bmatrix}$$

What is the value of the game in each case?

A payoff table of a two person zero sum game is given below.

$$A \begin{bmatrix}
10 & 7 \\
8 & 9
\end{bmatrix}$$

- a) Calculate maximin strategy of player A.
- b) Calculate minimax strategy of player B.
-) Is there a saddle point?
- 1) What proportion of the time should player A adopt his first and his second strategy?
- What percentage of the time should player B adopt his first and his second strategy?
- f) What will be the average value 'v' of the game?
- For the following games-
- 3) Reduce them to 2×2 matrices using the principle of dominance.
-) Hence determine the optimal strategies and the value of the game in each case.

A
$$\begin{bmatrix} 4 & 3 & 0 \\ -3 & -1 & -4 \\ 2 & 7 & 4 \end{bmatrix}$$
A
$$\begin{bmatrix} 5 & 4 & 11 \\ -1 & -10 & 0 \\ 2 & 3 & -11 \end{bmatrix}$$

bolve the following game after reducing it to a 2 × 2 game using dominance.

jolve the following game using the graphical method.

$$\begin{array}{c|cc}
 & B \\
 \hline
 -1 & 4 \\
 6 & 3 \\
 3 & 2 \\
 8 & 1
\end{array}$$

50he the following game using the graphical method.

A and B play a game of matching coins in which each has 3 coins; 10 paise, 25 paise players coins. Each player selects a coin without the knowledge of the other player, 25 paise of the coins is even, A wins B's coin and if the sum of coins is for Player A players A and B play a game selects a coin without the knowledge of the other player, 25 paise players coins. Each player selects a coin without the knowledge of the other player, 25 paise of the coins is even, A wins B's coin and if the sum of coins is odd, B wins A's choice. players coins. Each player A wins B's coin and if the sum of coins is odd, B wins A's coin. if the sum of anyoff matrix for Player A.

a) prepare a payoff matrix for Player A.

a) prepare a payoff matrix for Player A. of coins a payou man of coins apprepare a payou matches and the value of the game.

b) find the optimal strategy for each player and the value of the game.

A and B each taken. If only one of the players guesses correctly then the loser has to prove the players as is the correct answer, otherwise the game. A and B each take out only one of the players guesses correctly then the loser has to pay the opponent has taken. If only one of the players guesses correctly then the loser has to pay the A and b taken. It only matches to sponent has taken. It only many matches to pay the opponent has taken. It only many matches to pay the opponent as many rupees as is the correct answer, otherwise the game is a draw and the payout is obtain the payoff matrix. Num Obtain the payoff matrix. Formulate the following game as an LPP and obtain its solution.

formulate the tollowing game as an LPP and obtain its solution.

$$X \begin{bmatrix} 6 & 1 & 7 \\ 1 & 9 & 3 \end{bmatrix}$$

Y.B.Sc. Practical No.9, 1.1

VITAL STATISTICS

Calculate the crude and standardized death rates for the local Population from the following data e the crude death rate of the standard population.

and compa	Stand	lard	Local Local		
Age	Papulation	Deaths	Population	-	
Group	600	18	400	Deaths	
0-10	1000	5	1500	16	
10 - 20 20 - 60	3000	24	2400	24	
60 - 100	400	20	700	24 21	
60 - 10	TD D for Co	menters A on	41 1	41	

60-100 1 Annual on the basis of the following data.

Age (in yrs)	Specific dea country A (at rates Tol.	Life table population of	stationary of continent
	Males	Females	(stand	dard)
0-4	20.0	5.0	600	Females
5 – 14	1.0	0.5		700
15 – 24	1.4	1.0	2100	2000
25 – 34	2.0	1.0	5500	5150
35 – 44	3.3	2.0	4800	4650
45 – 54	7.0	5.0	4300	4100
55 - 64	15.0	12.0	4500	4200
65 - 74	40.0	35.0	4000	4300
75 & above	120.0	110.0	2600	3050
he number of bi	rths occurring in		1600	1850

The number of births occurring in a country in a particular year is shown here classified according to age of mother, together with the female population in each age group of the reproductive period.

			population in c
	Age	Female	Number of births to
	Group	population	mothers in the age- group
	15 - 19	84796	2349
	20 - 24	70018	14547
	25 - 29	72660	16746
	30 - 34	75924	10229
	35 - 39	75109	5257
	40 - 44	71625	
	45 - 49	66660	1432
•		ulation of the	93

The total population of the country during the year was 2, 285,800. With the above data, determine

b) the general fertility rate.

c) the age-specific fertility rates.

d) the total fertility rate.

(computed on the basis of females alone) are shown in the following table along

ates (tes.	Mortality rate	
The fertility rates (each. Tates. Fertility rate Fertility rate (per the cusand) 100 (per the cusand) 13.5 55.968	(per thousand)	
fertility Tablety Tertility Tand) The die mortality Tertility Tand) (per the usand) (per the usand)	0.40175	
The the mount per the table Group 13.5	0.42418	
Will Age Co. 55.968	0.45215	
15.20 70.2864	0.49320	١
	0.54330	1
1 7, 3,7	0.59699	1
30 70 15.2750	0.70473	
1.92	· Calcara data	_

45.50 NRR on the basis of above data.
45.50 Reproduction Rate (GRR) and 37 Rand NRR on the ball and Net Reproduction Rate (NRR) from the following Reproduction Rate (GRR) and Net Reproduction Rate (NRR) from the following

Compute Gross I	Female Population	Female births	Female Mortality rate
Group	(in '000s)	19423	0.189
	1478	70965	0.105
15-19	1604	95324	0.119
20 - 24	1778	62178	0.127
25 - 25	1687	33289	0.146
	1459	8713	0.162
35 - 39		608	0.189
40-44	1371	608	0.189

from the following data, calculate

Female Population	No. of births	Years lived
(in '000s)		by 1000
	· · · · · · · · · · · · · · · · · · ·	Female births
152.7	2971	4813
	22773	4790
	28245	4761
177.6	19548	4726
185.2	11462	4685
167.1	3235	4632
150.3	217	4552
	185.2 167.1	152.7 2971 164.2 22773 176.1 28245 177.6 19548 185.2 11462 167.1 3235

Assume that sex ratio at birth was 13 males to 12 females.

practica					
Complete the following table Age(x) 93102	e				
the follows	d _x	1000q x	L,		
aplete (x) Ex	-	0.62		T _x	c
compleage(x) 93102	-	0.66	32469313		
10 -	_	0.72			
11 .	-	0.80			
12 -	-	0.90			
13 -		1.00		•	
14 -	niù ma	1.12		٠,	
15	1.00	1.23		A TOTAL STREET	
16		1.33		-	
17	1	1.40		4.045	
18. 19 following table complete the following ℓ_x Age (x) 1273	-	1.40		4,842,446	
19 following table			91-10000 21-11	Addition from the same	
elete the Tone	d ×	$q_x = 1$	T_x	e o	
Complete (x) 1273	*	*	* *	*	
0 1129	*	*	*	*	
1 1000	*	*	* 58510	*	
1 1090 2 1066	*	*	* 57432	*	
2 1066 3 1049	_		- 56374	_	

Given that the probability that a life aged 30 will survive 10 years is 0.85 and the probability that a given that the probability that a given that the probability that of three persons." Given that the product 10 years is 0.6, find the probability that, of three persons all aged 30, at the aged 40 will survive ages 40 and 50 and at least one will die after age 50 least one will die between ages 40 and 50 and at least one will die after age 50.

last one will use aged 30, 35, and 40 respectively. Find the probability that

Three persons dies before age 45 while the others survive to age 55.

b) at least one them attains age 65.

[use the LIC (1970-73) ultimate table]

famortality table is represented by the function.

$$\ell_{x} = 1000 \sqrt{100 - x}$$

a) the probability of a life surviving from birth to age 19.

h) the probability of a life aged 36 dying before age 51.

c) the probability of a life aged 64 surviving to age 91.

Alife aged 50 is subject to a constant force of mortality of 0.048790. Find the probability that he

will die between the ages of 70 and 80.

F.Y.B.Sc. Practical No. 9. 1.3

LIFE TABLES-II

 $\int_{0.000}^{100} \ell_{30} = 33839.37, \ \ell_{40} = 33542.311, \ \ell_{50} = 32669.855, \ \ell_{10} = 33839.37, \ \ell_{10} = 33542.311, \ \ell_{10} = 32669.855, \ \ell_{10}$ $11^{1/30}$ = 33839.37, ℓ_{40} = 33542.311, ℓ_{50} = 32669.855, ℓ_{60} = 30039.787; fit Given ℓ_{30} is law of mortality $\mu_{x} = A + Bc^{x}$. Given '30 Makeham's law of mortality $\mu_x = A + Bc^x$.

Makeham's law of mortality $\mu_x = A + Bc^x$.

Makeham's law of mortality is 50 lakhs. On the assumption that it is stationary the total population of a city is 50 lakhs. On the assumption that it is stationary the total population experiencing H^M mortality, find The total purpose and a long is 30 lakhs. C population experiencing H^M mortality, find population of persons who are account.

population experiences who are aged 55 years or more.

a) the number of persons who are aged 25 years. the number of persons who are aged 25 years or more but less than 45 years.

b) the number of persons who are less than 18 years of age. b) the number of persons who are less than 18 years of age.
c) ine number of persons who are aged 70 years

c) the number of persons who are aged 70 years or more.

a) the number of a city is 95 lakhs. Assuming the number of a city is 95 lakhs. the number of population of a city is 95 lakhs. Assuming that it is a stationary population the population of a city, find the number of schools and collection in the population of a city is 95 lakhs. Assuming that it is a stationary population the population of schools and collection in the population of schools and collection of the population of schools and collection of the population of the population of schools and collection of the population of the p The population mortality, find the number of schools and college the city needs on experiencing H^M mortality, find the number of schools and college the city needs on experiencing of one school for every 1500 students and one college for experiencing 11 and college the city nee experiencing of one school for every 1500 students and one college for every 8000 the basis of one school for every 8000 and college for every 8000 and college for every 8000 the basis of one school for every 1500 students and one college for every 8000 the basis of one school for every 1500 students and college for every 8000 the basis of one school for every 1500 students and one college for every 8000 the basis of one school for every 1500 students and one college for every 8000 the basis of one school for every 1500 students and one college for every 8000 the basis of one school for every 1500 students and one college for every 8000 the basis of one school for every 8000 the basis of one school for every 8000 the residents in the city who are school for every 8000 the basis of one the basis of one some that 75% of the residents in the city who are aged 5 years or more students. Assume that 75% of study at school and 65% of the residents of the sound of students. Assume and the study at school and 65% of the residents, who are but less than 15 years or more but less than 20 years qualify for study at school and 65% of the residents, who are but less man 10 years or more but less than 20 years qualify for study at college.

aged 1) years of stable (period of selection 2 years) is given below find

a) 2 P[30] +1. b) 2 9 [31]+1

Age at entry	l [x]	l[x]+1	ℓ x+2	Age attained $x + 2$
[X]	486024	485094	484096	32
30	485028		483059	33
	484012		481994	34

ha certain select mortality table the period during which selection is assumed to have effect an the mortality experienced is four years. Given the following table and that $\ell_{[30]} = 494$, 985, find the values of

a) [120]+2 b) l 24

0) [21]

0/12/1

Age at entry	d _[x]	$d_{[x]+1}$	d[x]+2	d[x]+3	d[x]+4	Age-attained
[x]	,	[].				x + 4
20	641	778	864	907	927	24
21	640	781	866	913	933	25
22	640	782	871	919	942	26
23	639	785	876	928	950	27
24	639	789	884	936	963	28

T.Y.P.Sc. Practical No.9, 1, 4

ANNUITIES-I

A sum of Rs. 50 is contributed by a 25 years old person to his provident fund during the current month, what will this particular contribution amount to by the time he retires at the age of 60? Assuming a rate of interest of 8% p.a.? The compound interest on Rs. 1000 in four years at a certain rate is Rs. 200 and in 5

years it is Rs. 256 at the same rate. Find the rate of interest.

A sum of Rs. 100 is lent at interest of 12% p.a. convertible monthly. What is the total A sum of a sum of 12 years and 5 months? Calculate effective rate of interest.

Find the accumulated value at the end of 12 years and one and a half month of an amount of Rs. 750 invested at 9% p.a. for the first six years and thereafter at 10% p.a.

convertible quarterly.

A has a right to receive an amount of Rs. 500 at the end of 6 years from now. The right has been sold to B for a present value calculated at the rate of 7% p.a. The money thus received was invested by A in a deposit account at 9% p.a. payable half yearly. After 4 years the account had to be closed and A then invested the amount available at 5% p.a. How much has A gained or lost in the transaction, at the end of

6) Three loans of Rs. 800 each were given to Mr. Shah four years, three years and two years ago, respectively. The interest charged by the lender was 12% p.a. Mr. Shah could make a repayment of Rs.600 only at present with the promise to clear the dues at the end of three and a half years from now. How much will he have to pay then?

- 7) Three years ago A bought industrial share for Rs. 500 and has received on it three yearly dividends of Rs. 50/- each the first one having been paid one year after purchase. He wants to sell the share now and realize a rate of interest of 8% p.a. on the transaction. For how much should he sell?
- 8) A person has purchased a bond of the face value of Rs. 1000 on which interest is payable 7% p.a. He received in all three-interest payments, the first one falling due one year after purchase. At the end of three years if the bond is redeemable at no extra premium (i.e. on par) and if the person has realized an interest yield of 8% p.a. in the transaction, what is the purchase price?

Y.B.Sc. Practical No.9.1.5

ANNUITIES-II

The cash price of a scooter is Rs.40000/-. A company however offers installment The cash price of a scool and payment of Rs. 5000 is to be made and a series of 5 equal plan where an immediate payments made thereafter, the first installment being payable at the company wishes to realize a rate of incompany and the company wishes to realize a rate of incompany and the company wishes to realize a rate of incompany w the cash r an immediate representation and thereafter, the first installment being payable at the end representation, calculate half yearly months. If the company wishes to realize a rate of interest of 10% controlled months. plan who payments had been payable at the end half yearly in the transition, calculate half yearly installment? of six months. It the transition, calculate half yearly installment? half yearly in the transition of Rs. 200 p.a. for 10 when the transition of Rs. 200 p

of six in the nanisment? Convertible was payment of Rs. 200 p.a. for 10 years, the first payment being Mr. X is entitled to a payment is ceased at the end of 10 the payment is increased. Mr. V is entitled to a payment after ten years. The annual payment is increased to made one year from now and after ten years. The annual payment is increased to made one a, then the payment is ceased at the end of 15th year. Calculate the Mr. X arrate one year from now made one year from the payment is ceased at the end of 15th year. Calculate the percent ps. 400 p.a. then the payment is ceased at the end of 15th year. Calculate the percent ps. 400 p.a. and accumulated value at a computated value a Rs. 400 p.a. then the payments to Mr. X at rate of interest 7%p.a.? value and accumulated value at an effective rate of 9% p.a.? Calculate the present and accumulated value at an effective rate of 9% p.a. of an end of Rs. 120 p.a. payable quarterly for twelve years?

Calculate the present value and accumulated value are receive rate annuity due of Rs. 120 p.a. payable quarterly for twelve years? annuity due of Rs. 12 represent value and accumulated value at the rate of interest of 7% p.a. Calculate the present under which 10 payments of Rs. 600 each are made at the of a deferred and deferred and deferred and deferred of every three years, the first payment being due 20 years hence.

obtain the present value of an increasing annuity for ten years under which the first Obtain the present of Rs. 200 is paid one year from now and the successive payments increase payments are year. Assume 8% rate of interest. payments of per year. Assume 8% rate of interest.

by Rs. 150 per year.

Appropriate a loan of Rs. 2 lacs at 10% rate of interest for house repair which he has to repay in 20 equal installments. Find

a) annual installment

b) the interest contained in the eighth payment

c) principal outstanding after the 8th payment is made.

Assume that first payment is made at the end of one year from now.

Calculate the present value of an

a) immediate perpetuity and

b) perpetuity due

The payment is done at an effective rate of interest of 8% p.a. of Rs. 180 p.a. payable twice a year.

Calculate the present value of an immediate perpetuity of Rs.200 p.a. payable half yearly at a rate of interest of 6% p.a. convertible half yearly. The first payment being received at the end of six years from now.

Y.B.Sc. Prictical No.9.1.6 PREMIUMS AND ASSURANCES

Consider the following data. 43 44 45 79999 78942 79488 78357 80480 80935

ds 455 401 405 626

ds 455 (alculate, assuming rate of Rs. 5000/- for 3 years for a person aged 40 years, a) temporary assurance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 42 years received and the surance of Rs. 3000/- for a person aged 40 years received and the surance of Rs. 3000/- for a person age (3) temporary assurance of Rs.3000/- for a person aged 40 years.

1) temporary assurance of Rs.3000/- for a person aged 42 years receivable on the endowment assurance of Rs.3000/- for a person aged 42 years receivable on the endowment age 45

attaining age 45

years.

years.

c) double endowment assurance of Rs. 10000/- for 3 years for a person aged 42 years.

c) double emporary assurance of Rs. 10000/- for 3 years for a person aged 42 years. c) double endowment assurance of Rs. 10000/- for 3 years for a person aged 42 years for a person aged 40 increasing temporary assurance of Rs. 10000/-

years.
e) deferred temporary assurance of Rs.10000/- on a person aged 40 years for 3 years

deferment period of 2 years. Using commutation functions based on LIC (1970-73) ultimate mortality @ 6%

interest, calculate

for a person aged 30, the present value of for a period of Rs. 15000/- for a period of 25 years.

b) whole life assurance of Rs. 50000/-

c) increasing whole life assurances of Rs.5000/-

occurs after attaining age 35 years.

c) special endowment assurance providing both increasing death benefits and increasing survival

benefits of Rs.5000/- for a period of 10 years.

An endowment assurance of Rs. 10000/- is given to a person aged 28 for a terms of 10 years with

the following benefits.

a) one fifth of the sum assured at the end of every alternate year in case the person survives till that

period

b) full sum assured at the end of year of death, whenever it occurs during the term of 10 years.

find the value of this assurance.

Basis: LIC (1970-73) table at 6 %.]

on the basis of the LIC (1970-73) ultimate table at 6 %, calculate the net level on the basis of the Lacon assured of Rs. 10000/- for the following assurance on annual premiums for a sum assured of Rs. 10000/- for the following assurance on

a) whole life assurance.

a) whole life assurance, premiums limited to 20 years.
b) whole life assurance for 25 years, premium downent assurance for 25 years.

b) whole life assurance for 25 years, premiums limited to 15 years.
c) endowment assurance -the assurance to commence of the c b) with the continue assurance of the assurance to commence at age 35 years and the continue

then to continue

for 10 3000 e) double endowment assurance for 25 years. e) double the net level annual premium under a special Endowment assurance of Rs. Calculate the net level annual premium under a special Endowment assurance of Rs. Calculate the first 10 years. In the event of death during the first 10 years total 30000/- on (30) for returnable and in the event of death during the last 10 years total premiums paid are returnable and in the event of death during the last 15 years or returnable to the end of 25 years, the basic sum assured becomes premiums paid at 25 years, the basic sum assured becomes payable. [Basis: UC (1970-73) and 6 % interest.]

A special policy provides for the following benefits. A special policy of Rs. 1000/- with guaranteed annual additions of Rs. 400/- for each

premium paid after the first, if death occurs within the term of assurance, b) Rs.10000/- payable on survivance to the end of term of assurance and b) Rs. 10000 Positive and b) Rs. 20000/- at death after expiry of the term of assurance.

Calculate net

annual premium under the policy on the life aged 30 years for a term of 25 years. [Basis: LIC (1970-73) table & 6% rate of interest.]

A life aged 30 effected an endowment assurance policy of Rs. 10000/- payable at the end of 25 years or at previous death, with the condition that the premiums payable during the first 5 years shall be ordinary whole life premium, the premium payable thereafter being correspondingly increased. Calculate the annual premium during the last 20 years using LIC (1970-73) tables at 6% rate of interest.

1) Alife aged 40 effected an endowment assurance policy of Rs. 5000/- payable at the end of 15 years or at previous death with the condition that the annual premium payable during the first 5 years is three times the annual premium payable thereafter. Calculate the annual premium payable during the first 5 years using LIC (1970-73) ultimate table at 6% rate of interest.

9 Calculate net annual premium under a special endowment assurance of Rs. 18000/on (35) for 25 years, the premium payment limited to 20 years. In the event of death during the term of assurance, total premiums paid are returnable and on survivance to the end of 25 years, the basic sum assured becomes payable. [Basis: LIC (1970-73) table.]

T.Y.B.Sc. Practical No.9.1.7 LIFE ANNUITIES AND TEMPORARY ANNUITIES ing data

	the follo	owing da	ta.	1	Art desi				
	Consider the follo	50	51	52	53				
11	- 111	01077	91261	'90484	89622	54	55	56	_
	No. of persons	32. 8 6	. D.10.	3.7.11,21	03032	88703	87689	20	57
			interest	'90484				86282	85387
	living (2 x) Assuming 6% p.a	annuity o	f Rs 100	000/	, using al	ove data	the		
	assumptiary life	almuny o	1 100.100	/00/- p.a.	for a ne	TEOn -	o are pre	sent valu	200

Assuming 676 P. Assuming 676 P

first payment being made immediately.

first payment to be first

years, the first payment being made on attaining age 53 years.

years, the his property life annuity due of Rs.2000/- p.a. to a person aged 50 years for 3 period of 4

years with a deferment period of 2 years.

years what years who had you what you who had you what you what you want you what you want you want you what you want y 6% p.a. rate of interest compute the present value of

a) an immediate life annuity of Rs. 10000/- p.a. for a person aged 60 years.

a) an infinited an all all an annuity of Rs. 10000/- p.a. for 10 years to a person aged 50 years.

c) a life annuity due of Rs. 10000/- p.a. to a person aged 50 years.

d) a deferred temporary immediate life annuity of Rs. 10000/-p.a. for 20 years to a person aged 40

years; deferment period being 5 years.

c) an increasing temporary life annuity due of Rs. 1000/- p.a. for 25 years to a person aged 30 years.

3) A person received a bonus of Rs.60000/- at the time of his retirement (i.e. at age 60 years). He wishes to buy an increasing life annuity with Rs. 5000/- as the first payment payable at the age 62 years and the amount increasing by Rs.A every year. Find the value of A based on the commutation functions of LIC (1970-73) ultimate

4) A person wishes to keep some amount with a trustee firm so as to provide a life annuity of Rs.2500/- p.a. to his grand son, now aged exactly 18 years, the first payment to be made on his 25th birthday. How much amount of money should he deposit today with the firm so as to provide the desired annuity to his grand son? [Use LIC (1970-73) ultimate mortality tables.]

An annuity due of Rs. 200/- p.a. is payable to a life aged 60 years. If the annuitant wishes to alter the annuity into an annuity commencing Rs.100/- and increasing by a

level amount each year, determine the amount of annual increase.

[Basis: LIC (1970-73) ultimate table using 6% p.a. rate of interest]. Find the numbers living at ages 25 to 28 years if at 6%


 $a_{25} = 15.05$, $a_{26} = 14.974$, $a_{27} = 14.892$, $a_{28} = 14.806$, $a_{29} = 14.715$, $\ell_{29} = 98206$. Find the value of a pure endowment benefit of Rs.6000/- for a person aged 40 years

-- 12 060 and a 40. Payable on his surviving to age 60 at 6% p.a. given that $a_{40:207} = 13.060$ and $a_{40:10} = 13.060$ and $a_{40:207} = 13.060$ and $a_{40:207} = 13.060$


T.Y.B.Sc. Practical No.9.2.1

RELIABILITY

A system consists of 10 components A,B,C,D,E,F,G,H,I and J with reliabilities 0.8, 0.75, 0.99, 0.95, 0.85, 0.98, 0.7, 0.88, 0.96 and 0.95 respectively which are connected 0.75, 0.99, in the diagram given below. Find the reliability of the system.



2) Find the reliability of the configuration given in the following figure:



(Note: The figures indicate reliabilities)

3) A set of tests can be performed to detect a disease before it develops fully and the condition goes beyond cure. But none of these tests are confirmatory in nature. The sets have series of stages to be performed sequentially. The tests are denoted by Ai (i = 1, 2...9) and Bj (j = 1, 2...9). Stages and tests along with their probabilities of failure are listed below.

Set A

- (i) A₁ (0.05)
- (ii) either $(A_2 (0.10) \text{ and } A_3 (0.10) \text{ together})$ or $A_4 (0.15)$
- $(iii) A_5 (0.12)$
- (iv) either A_6 (0.15) or A_7 (0.12) or A_8 (0.05) and A_9 (0.05) together.

Set B

- (i) either B₁ (0.05) or B₂ (0.07)
- (ii) B_3 (0.12)
- (iii) either B_4 (0.08) and B_5 (0.15) together or B6 (0.09) and B_7 (0.20) together. Determine the reliability of each of the two sets of tests.

An electronic circuit consists of 15 valves, 20 resistors and 10 capacitators, all An electronic circuit Components in each category are identical and their failure connected in series. The componential distribution with the following An expense of the series of th

Mean Failure Time (in hours) 10,000

time: Components Vaives 20,000 Resistors 20,000.

Capacitators What is the reliability of the system for 100 hours? Also, find the MTBF of the system.

- Suppose the failure time of a component is exponentially distributed. Ten identical suppose the failure times in hours are observed to be as follows: components $t_1 = 480$, $t_2 = 630$, $t_3 = 560$, $t_4 = 500$, $t_5 = 495$. $t_6 = 475$, $t_7 = 562$, $t_8 = 478$, $t_9 = 614$, $t_{10} = 566$. t₁=480, t₂ Estimate the hazard rate and the reliability of the component for 500 hours.
- of An equipment consists of 100 parts distributed in 3 branches A (20), B (20) and C (60). A is connected in series with a branch which has B and C connected in parallel. In each branch, the parts are themselves connected in series. Each part has constant hazard rate 0.005. Compute the reliability of the equipment at 10 hours.

TIME SERIES

Practical No.9.2.2 The following data show the number of thefts (in thousands) in the United States for the legislation of the original data. ollowing data show and trend line using the method of semi-averages for the data.

the years the graph of Also, plot the 1985 1986 Year 19.0 20.6	1987 1988	1989	1990	1991	1992	1993	1994	1995
Year 1900 20.6	20.1 20.7	27.5	2.7.4	24.7	23.8	24.5	23.3	21.6
10.0f	-f thefte	for the ve	ar 1990	6				

thefts predict the number of thefts for the year 1996. the predict the number of the sales (in thousand Rs.), determine the trend values ming a parabolic trend in the sales (in thousand Rs.)

Assuming data.	1994	1995	1996	1997
for 110 1991 1992 13	20	19	17	15
7 1 1 2 1	or 1998			

Also, estimate sales for the year 1998.

Also, established the trend values for the following data by the method of least squares is Calculate the trend. assuming a linear trend.

Year	Quarter						
1 cai	I	п	Ш	IV			
1995	25	30	21	32			
1996	27	28	25	34			
1997	22	27	21	30			
1998	24	25	20	33			

(alculate seasonal indices by the ratio to moving average method for the following

Year	Quarter						
1 cal	I	П	m	IV			
1981	68	62	61	63			
1982	65	58	66	61			
1983	68	63	63	67			

Using ratio to trend method, compute seasonal indices for the following data.

Quarter	Year						
- Cuartor	1980	1981	1982	1983			
I	75	86	90	100			
П	60	65	72	78			
III	54	63	66	72			
IV	59	80	85	93%			

Calculate seasonal i			Ou	arter	
Calcul	Year	I	п	III	IV
	1991	6.0	6.5	7.8	8.7
	1992	5.4	7.9	8.4	7.3
	1993	6.8	6.5	9.3	6.4
	1994	7.2	5.8	7.5	8.5

SIMULATION

Trucks arrive at a certain toll post according to the following frequency distribution.

4 12 27 23 2.5 3.0 3.5 Time between arrivals (in min) 3.0 3.5 12 27 23 16

frequency frequency frequency with 0.5 1.0 1.5 2.0

distribution. Service time (in min) 1.0 0.51.5 2.0 35 27 25 13

Frequency freque Estimates by simulating for 10 arrivals.

server by summary server by su

for arrival distribution: 74, 29, 92, 62, 31, 44, 24, 70, 52, 43 for service distribution at time t = 0. First customer arrives let for service and for service and for service at time t = 0. First customer arrives later.]

[Observation begins at time t = 0. First customer arrives later.]

[Observation of a dealer for a certain product, the probability distributions of the daily demand and the

lead time are as follows.

Daily Demand Distribution

10			July L	cinani	DISH	<u>iounor</u>	1			
a mand	3	4	5_	6	7	8	9	10	11	_
Demand Probability	0.02	0.08	0.11	0.16	0.19	0.13	0.10	0.08 ($\frac{11}{107}$ 12	1
1700401117		<u> </u>	Lead T	ime Di	stribut	ion	3.10	0.00	0.07	6
	()	2	2	4	-	1				

Lead Time (days) 0.20 | 0.30 | 0.35 | 0.15 P:obability '

Shortage cost (C₂) is Rs.20 per unit per day, ordering cost (C₃) is Rs.80 per order, holding cost (C₁) is Rs.2 per unit per day, opening stock on 1st day is 30 units. Consider a simulation plan which calls for a reorder quantity of 40 units and reorder level of 20 units. It is assumed that orders are placed / received before the end of the day. Using following random numbers generate demand for 15 days and by simulation arrive at average cost per day.

Random Numbers (Demand): 68, 13, 09, 20, 73, 07, 92, 99, 93, 18, 24, 22, 07, 29, 57.

Random Numbers (Lead time): 47, 74, 25, 21, 47

A Company manufactures 30 units per day. The sale of these items depends upon demand which has the following distribution.

Demand (units) 27 28 29 32 30 31 Probability 0.10 0.15 | 0.20 0.35 | 0.15 | 0.05

The production cost and sale price of each unit are Rs.40 and Rs.50, respectively. Any unsold product is to be disposed off at a loss of Rs.15 per unit. There is a penalty of Rs.5 per unit if demand is not met. Using the following random numbers Randon the total profit/loss for the company for the next ten days.

Genardom Numbers 10, 99, 65, 99, 95, 01, 79, 11, 16, 20. Generate Random observations from each of the following (5 in a, b and 2 in c).

Uniform distribution: $f(x) = \frac{1}{18}$; 6 < x < 24

= 0; otherwise

Section with mean = 2. $\sqrt{\frac{1}{2}}$ distribution with m = 3, $\lambda = 0.4$

of a factory is normally distributed with mean $\mu_1 = 200 \text{ kg}$ and the daily production for a factory is normally distributed with mean $\mu_1 = 200 \text{ kg}$ and $\mu_1 = 200 \text{ kg}$ and $\mu_2 = 20 \text{ kg}$ and $\mu_3 = 20 \text{ kg}$ and $\mu_4 = 200 \text{ kg}$ and The no. of production hours is also normally distributed with mean $\mu_1 = 200 \text{ kg}$ and $\mu_1 = 200 \text{ kg}$ and $\mu_2 = 200 \text{ kg}$. Simulate the daily production and daily production. The no. of production is also normally distributed with mean $\mu_1 = 200 \text{ kg}$ and $\mu_1 = 200 \text{ kg}$ and $\mu_2 = 20 \text{ kg}$. Simulate the daily production and daily production $\mu_1 = 200 \text{ kg}$ and $\mu_2 = 200 \text{ kg}$ and $\mu_3 = 200 \text{ kg}$ and $\mu_4 = 200 \text{ kg}$ and $\mu_5 = 200 \text{ kg}$ and $\mu_6 = 200 \text{ kg}$ and μ_6 sides and s.des to estimate the daily production rate. Use the following random for ten days to estimate the daily production rate. The following random for ten days to estimate the daily production rate. The following random for ten days to estimate the daily production rate. The following random for ten days to estimate the daily production rate. The following random for the following random rate.

pumpoduction: 62, 65, 20, 95, 59, 21, 84, 87, 39, 87 for production hours: 62, 65, 20, 95, 59, 21, 84, 87, 39, 87

dentist schedules all her patients for 30-minute appointments. The following schedules all her parameters of work and the probability distribution of

A mary shows amplete 1	he work.	dis dis
summary shows the summary show	Time Required (minutes) 45	Probability
and the ory of Deman	45	0.40
Filling	60	0.40
Crown	15	0.15
:00		0.13

Cleaning

Extraction

Check-up

Assuming that all patients arrive at the clinic at their scheduled arrival time, starting Assuming that an personal state of the dentist's clinic for 7 patients and determine average waiting ine for the patients as well as idle time for the dentist. Use the following random numbers: 40, 82, 11, 34, 25, 66, and 17.

45

15

0.10

0.20

```
T.Y.B.Sc. Practical No.9.2.4
                 MATHEMATICAL ECONOMICS- I
```

The demand function at point X = 4 and P = 6 is given by X = 10 - P. If the price the demand and hence $\frac{1}{2}$ the electricity of demand $\frac{1}{2}$ The demand function to the relative decrease in demand and hence an increases by 5 %, determine the relative decrease in demand and hence an increase in the result with the description to the clasticity of demand. Compare the result with the simulation to the classicity of demand. The $\frac{1}{100}$ section of the elasticity of demand. Compare the result with the correct value approximation of demand corresponding to X = 4. approximation of demand corresponding to X = 4.

of clasticity of demand and price of the sand P_1 are demand and price of the sand P_2 are demand and price of the sand P_3 are demand and price of the sand P_4 are demand and P_4 are demand P_4 are demand and P_4 are demand and application of definition of the state of t If X₁ and the demand functions are given by coffee and the p-1.3 p.0.5

 $X_1 = P_1^{-1.3} P_2^{-0.5}$ $X_2 = P_1^{0.3} P_2^{-0.5}$

N2 - 1 Show that two commodities are competitive. Also obtain four elasticities of demand. Given the demand function

 $p = 100 - 0.5 \text{ X}^2$ and the cost function

$$P = 100 - 0.3 \text{ }^{2}$$

$$C = \frac{X^{3}}{3} - 6 X^{2} + 108 X,$$

determine a) the equilibrium output.

a) the equalities and profit after the imposition of a tax of Re. 1 per unit b) the change in price, output and profit after the imposition of a tax of Re. 1 per unit

of output.
c) the price if the government grants subsidy of Re.1 per output.

c) the price \mathbb{R}^2 in Polymer output.

Amonopolist produces amounts X_1 and X_2 of two goods A and B at the total cost \mathbb{R}^2 in $C = X_1^2 + 4X_1X_2 + 6X_2^2.$

The demand functions for these goods are $P_1 = 40 - X_1$ and $P_2 = 100 - 5 X_2$ respectively, where P₁ and P₂ are the prices. Determine the quantities and prices, which maximize the total profit. Also find maximum profit.

The demand functions of discriminating monopolist are $P_1 = 12 - q_1$ and $P_2 = 20 - 3$ q_2 and the total cost function is $C = 3 + 2 (q_1 + q_2)$. Determine the prices set and quantities sold in the two markets that will maximize the total profit. Show that discriminating monopolist charges more price in the less elastic market.

Assume that the demand and cost function of daopolisis are

$$P_1 = 100 - 2q_1 - q_2$$
, $C_1 = 30 q_1$ and

$$P_2 = 95 - q_1 - 3q_2$$
, $C_2 = 25 q_2$ respectively.

Find the optimum prices, quantities and profits when

a) duopolists recognize their dependence.

b) duopolists do not recognize their dependence.

The cost function of a firm is given by $C = 2 Q - 2 Q^2 + Q^3$.

a) AC function

h) MC function

c) at what level of output AC is minimum?

(1) Verify that at minimum average cost AC = MC.

Given the demand function P = 86 - 25 X, verify the relationship between marginal texture. revenue, average revenue and elasticity of demand (η_d).

T.Y.B.Sc. Practical No.9.2.5 MATHEMATICAL ECONOMICS- II

Given the production function $Q = f(L, K) = LK - 0.2 L^2 - 0.8 K^2$. Construct the given the productivities for labour (L). Assuming that K = 10Given the production productivities for labour (L). Assuming that K = 10, examine average and marginal production function. the three stages of the production function.

the three stages of the production is given by $Q = 5 L^{0.3} K^{0.7}$, where L and K represent the production capital respectively. The prices per unit of labour and capital the production and capital respectively. The prices per unit of labour and capital are Rs.20/-labour and/r respectively. If the cost is fixed at Rs.500/-, determine the labour and capital are Rs.20/- respectively. If the cost is fixed at Rs.500/-, determine the amounts of and Rs.40/- respectively and capital in order to maximise the output. and Rs. 40/2 and capital in order to maximise the output.

About and capital in order to maximise the output.

A production function is O = 5 T 0.777 h 3.

A firm's production function is $Q = 5 L^{0.7} K^{0.3}$. The price of labour is Re.1 per A firm's price of capital Rs.2 per unit. Find the minimum cost A firm's production of capital Rs.2 per unit. Find the minimum cost combination of unit and labour for an output rate of 20. capital and labour for an output rate of 20.

The firm's production function is

$$q = 12 - \frac{L + K}{L.K}$$

The prices of labour, capital and output are Rs.1, 4 and 9 respectively. Find the maximum profit combination of capital, labour and output.

INPUT-OUTPUT MODELS

Practical No.9.2.6 Assume that the economy is divided into two sectors and input output table in Assume mits is as follows.

Is is	Purchasir	Final	
Producing Sectors	I	II	Demand
and the second second second second	200	250	50
17	100	500	400

Obtain the technology matrix. Obtain the recumerous of the two sectors are desired to be 180 and 440 units for b) If the final demands of the total output in

sectors I and II

respectively, then find the total output in accordance. respectively, then \underline{F} which is consistent with the output vector \underline{X} and \underline{F} ind the demand vector \underline{F} which is consistent with the output vector \underline{X} and \underline{F} independent \underline{A} . echnology matrix A.

$$\underline{X} = \begin{bmatrix} 25 \\ 21 \\ 18 \end{bmatrix} \qquad ; \qquad A = \begin{bmatrix} 0.2 & 0.3 & 0.2 \\ 0.4 & 0.1 & 0.2 \\ 0.1 & 0.3 & 0.2 \end{bmatrix}$$

Compute the gross value added in each sector and show that the total value added in tach sector and total final demand is equal.

A three sector input-output matrix [I - A] is given as

$$\begin{bmatrix} 1 & -0.5 & 0 \\ -0.2 & 1 & -0.5 \\ -0.4 & 0 & 1 \end{bmatrix}$$

With labour coefficients (0.4, 0.7, 1.2) and the household demand for the outputs of three sectors as 1000, 5000 and 4000 respectively, determine the level of output. If the wage rate is Rs. 10/- per labour day, find the equilibrium prices of the commodities in the three sectors.

Assume that the economy is divided into three sectors and input-output table in physical units is as follows.

Producing	Purcl	hasing Se	Final	
Sectors	I	II	Ш	Demand
I	100	150	225	.75
П	135	150	300	115
III	270	200	300	130
Labour (in lacs)	12	10	16	1/2

the final demands for the products of the sectors I, II and III rise to 100, 200 and M respectively, find the corresponding rise in the employment.

水水水水水水水水水水水水水水水水水水水水水水水水水